IMPACT OF H5N1 INFLUENZA ON DAIRY COWS

Arifa MEHREEN

Muhammad MAJEEBULLAH

Shagufta ABBAS

Muhammad SAFDAR

1. Introduction

The H5N1 flu virus, commonly referred to as avian flu or bird flu, is one of the most threatening diseases to animals and humans worldwide in relation to agriculture and food sector, especially dairy (Duarte et al., 2024). Although the H5N1 strain of the avian influenza virus is mainly reported in birds, its zoonotic nature makes it infect mammals, particularly dairy cows, which has elicited a lot of concern among veterinarian professionals, farmers and researchers. H5N1 threat in dairy cows is complex since it adds multiple issues concerning animal health, milk production and socio-economic situations. H5N1 was first reported in birds in the late twentieth century and has been responsible for several outbreaks fowl resulting in a high mortality in poultry population (Shi et al., 2023). Described as a zoonotic virus, hence one capable of spreading from birds to humans and other animals, the virus has received significant attention from researchers. It is therefore necessary to understand the means through which H5N1 spreads to affect dairy cows. Transmission involves direct contact with infected bird and its excretion feces, feed and water, and other ways like contaminated equipment and through humans(Smith et al., 2020). Wild birds, especially birds that migrate, usually possess the virus and can spread it to other areas ad farms. These numerous pathways of transmission reinforce the importance of sound animal health measures and supervision. Avian flu H5N1 can be devastating to dairy cows, causing illness and potential direct or indirect reduction in production. In particular, affected cows may demonstrate symptoms like respiratory disorders and gastroenteritis as well as severe neurological signs. These symptoms can indeed lead to decreased milk production and poor quality thus being a major concern to most farmers who rear dairy animals. Also, the infection with H5N1 of dairy herds leads to the ban of exports and the decline of consumer confidence, which only increases the losses (Gulati & Juneja, 2023).

Prevention of H5N1 in dairy cows relies heavily on early identification and passive control. The identification of early clinical manifestations alongside testing for the virus assists in quarantine and entails restrictions of the virus transmission. Measures such as supportive therapy for effected animals, implementation of disease control measures and disinfection, as well as the use of vaccines where applicable. The social and economic consequences of H5N1 in dairy populations range extensively. In addition to the direct monetary costs, outbreaks can interfere with local and global dairy production, leading to scarcities and price shifts in food products. The level of psychological disruption is also present in farming societies, which might be forced to slaughter their livestock, and lose jobs (Kappes et al., 2023).

2. Pathways of H5N1 Transmission to Dairy Cows

H5N1 is a virus that is pathogenic mainly to birds but occasionally affects dairy cows through direct and indirect transmission (Graziosi et al., 2024).

Direct Transmission: This occurs through the direct interaction with the bird or their excreta which include feces, saliva and nasal discharge (Ly, 2024). This can be through the cows ingesting the plants that have been contaminated or through the consumption of contaminated water by the animals. It raises the likelihood of transmission specifically where poultry and dairy farming is done in a combined manner as birds and other animals together with farming tools can easily spread the

virus. Direct Pathway of H5N1 transmission to dairy cows is shown in the Figure 1.




Figure 1 Direct Transmission of H5N1 Transmission to Dairy Cows

Indirect Transmission: It is also passaged by contact with things like tools, cloths, or vehicles bearing the stamp of the virus. This implies that the farm workers, veterinarians and visitors can carry the virus in their belongs from infected poultry farms to dairies. Indirect transmission of H5N1 to dairy cows is shown in Figure 2. It may also be transmitted indirectly through water as the virus can spread on the surface of water after being carried there by dead wild birds (Bauzile et al., 2022).




Figure 2 : Indirect Transmission of H5N1 Transmission to Dairy Cows

Airborne Transmission: While considered less probable, H5N1 can also spread through aerosols carrying infectious droplets. Avian influenza viruses are expelled by infected birds as aerosolized particles which can be inhaled by the dairy cows particularly when they are locked or when the houses are inadequately ventilated (Bhalerao et al., 2023).

Human-Mediated Transmission: Humans that are infected with the virus can pass the virus from one human to another, and in the same manner H5N1 can be transmitted from human to other animals. The infected poultry handlers spread the disease to other animals including the dairy cows either through direct contact or through contacting clothes or equipment that has traces

of the disease. There is also fear that through the use of vision and voice such items as feed and bedding materials can transmit the virus from one farm to another (Huang et al., 2023).

Wildlife Reservoirs: Wild mosquitoes infected with H5N1 can fly far and spread the virus across regions and possibly to new regions such as the dairy farms (Marrana, 2022).

Contaminated Feed and Water: Feeding and watering are other means through which H5N1 migrates from one poultry house to another since the feed and water sources are also contaminated (Beerens et al., 2021). Virus dissemination is possible through droppings of infected birds and water sources can serve as a vector for the virus. The food and water that is assumed to have been contaminated with H5N1 must always be clean and safe for use.

3. Clinical Manifestations of H5N1 in Dairy Cows

In dairy cows, the clinical signs of H5N1 influenza are unspecific and affect several systems in the body. The initial clinical signs, though often vague, may be fever, anorexia, and nasal discharge. In later stages of the infection, the respiratory signs are characterized by persistent coughing, dyspnea and nasal and ocular discharge (Rosone et al., 2023). Some of the cows may also develop symptoms such as diarrhea and stomach ache. In more serious cases, neurological complications like ataxia, tremor, seizures are observed.

H5N1 also affects the quantity and standard of the milk produced by the animal. Mastitis in cows leads to reduced milk yield and the quality of milk is affected with higher somatic cell counts and can be tampered with by bacterial infection (Yoo et al., 2022). In severe cases, blood or clots can be observed in the milk being secreted by the mammary glands.

Autopsy of affected animals usually shows pathological changes in the lungs, trachea, bronchi, and gastrointestinal systems. In patients with neurological manifestations, neurological lesions may also be seen (Zappulli et al., 2020).

H5N1 infection in dairy cows can be identified through clinical and laboratory examination including PCR to identify H5N1 genetic material. Management mainly consists of supportive measures such as isolation, adequate fluid and electrolyte replacement, nutrition, and antibiotics for secondary bacterial infections. The outcome depends on the extent of the infection as well as the response to treatment.

4. Impact on Milk Production and Quality

H5N1 avian influenza infection in dairy cows leads to a significant decrease in milk production due to several factors: Focal infection, especially common in the form of fever and illness, dehydration, and respiratory troubles are some of the complications (Polley et al., 2022). These it depletes the energy, appetite, and water in the cow, which are all required to produce milk in large quantities. Likewise, (Kim et al., 2010) also consider that H5N1 has an effect on the physical distress of the affected animals, which in turn aids to divert energy from milk production.

Table 1: Impact on Milk Production and Quality

Impact Area	Observed Effects	Underlying Mechanisms
Animal Health	• Fever, illness, respiratory distress • Increased susceptibility to other diseases	• Direct pathogenic effects of H5N1 virus • Potential for secondary bacterial infections due to weakened immune system • Immunosuppression caused by H5N1 infection
Milk Production	Significant decrease in milk yield	• Focal infection (fever, illness) leading to energy depletion and reduced appetite • Dehydration and respiratory issues further hindering milk synthesis • Physical distress diverting energy from milk production (Kwon et al., 2005)
Milk Quality	Contamination of raw milk	• Presence of H5N1 virus particles or secondary bacterial contamination • Increased somatic cell count (SCC) indicating inflammation or mastitis • Altered fat and protein content due to stress and illness, impacting taste and processing properties
Economic Impact	Reduced profitability of dairy animal production	• Lower revenues due to decreased milk quantity and quality • Increased costs for veterinary services, feed, and potential animal removal • Negative market repercussions due to milk safety concerns, leading to lower milk prices or market restrictions • Long-term economic impacts: decreased returns on dairy farming, potential unsustainability for small farmers due to vaccination, culling, or movement restriction costs (especially with increased H5N1 circulation)

Apart from the raw milk production, H5N1 affects the quality of raw milk since infected cows have contaminated milk as shown in Table 1. The milk also becomes more inclined to exhibit increased somatic cell count (SCC) signifying inflammation or mastitis which hampers the quality of milk making it less preferable for processing. In aggravated cases, the milk may contain the virus particles or secondary bacterial contamination which may be hazardous to the health of consumers. They also lead to stress and illness, which can reduce milk production or increase the fat and protein content, resulting in poorer taste and prospectively and processing properties.

These changes in milk production and quality therefore have a direct impact on the economic profitability of dairy animal production. These are the animal productivity losses and include a decrease in the quantity and quality of milk produced, which may lower revenues while costs for veterinary services, feed and, in some cases, removal of affected animals rise. Furthermore, issues relating to milk safety or quality may have negative repercussions on markets and customers, pushing down milk prices, or even limits access to market by consumers. The more the H5N1 viruses circulate, the greater the potential for long-term economic impacts: the returns on dairy farming can decrease significantly and, for small farmers in particular, may not be sustainable due to costs associated with vaccinations and culling or movement restrictions (Kim et al., 2010).

5. Immune Response to H5N1 Infection

It has been observed that a dairy cow's immune system initiates a complex kind of defense

mechanism when it is infected with the H5N1 influenza virus (Alqazlan et al., 2022).

1. Innate Immune Response: This is the first response that the human body deploys in its defense and entails putting up barriers such as mucous membranes and recruitment of cells including macrophages and dendritic cells (Peiris et al., 2009). These cells help in preventing viral replication and also release special chemical substances called cytokines which help in regulating the overall immune response. It also stimulates the inflammation to confine the infections as part of the innate response.




Figure 3: Immune Response and Vaccination Strategies in dairy cows

2. Adaptive Immune Response: This response is more specific and entails the development of antibodies by B lymphocytes cells that can specifically identify and inactivate the H5N1 virus. Many other immune cells are involved in this fight, including specialized T cells: cytotoxic T lymphocytes that are able to find and kill infected cells. Immune Response and Vaccination Strategies in dairy cows are shown in figure 3. Once the pathogens infect the body, immunity cells are created as a memory that lasts and can protect a person in case of subsequent exposure to H5N1 (Lee et al., 2009).

Challenges In Immune Response

1. Antigenic Variation: The H5N1 virus demonstrates high mutation rates and antigenic drifts that alter the virus's surface proteins. This makes escapes the immune system and thus can easily produce new strains that will be very hard for the immune system to combat (Velkov et al., 2013).

2. Immunosuppression: H5N1 has immunosuppressive properties that weaken the immunity of the cow and causes susceptibility to other diseases (Schat & Skinner, 2022).

3. Variable Immune Responses: There are differences from one cow to the next in terms of its immune system and thus it is not the same affected or protected (Trevisi & Minuti, 2018).

Vaccination Strategies

It was found that vaccination remains a very effective option in the prevention and management of H5N1 infections in dairy cows. Strategies such as proper immunization can help increase the level of immunity in a population, limit the virus's circulation, and minimize the consequences of the epidemic (Abdelwhab & Hafez, 2012).

Types of Vaccines

1. Inactivated Vaccines: These consist of inactivated viruses that are able to trigger production of antibodies within the immune system besides not being able to cause an illness.

2. Live Attenuated Vaccines: These employs attenuated live viruses which are very effective in the production of immunity though they pose a slight risk of inflicting illness.

3. Subunit Vaccines: Of these, some contain viral proteins or antigens and are therefore safe but their effectiveness may need to be boosted through the addition of adjuvants.

4. Recombinant Vector Vaccines: These employ non-pathogenic viruses or bacteria which act as vectors to present the H5N1 antigens and elicit persistent immunity.

Vaccination Protocols

Effective vaccination protocols include (Gisbert et al., 2012):

1. Primary Vaccination: The first set of shots to help create immunity against a pathogen or toxin.

2. Booster Shots: A booster that every now and then is required to boost the immune system.

3. Herd Vaccination: This involves vaccinating all the animals within a given population to enhance the levels of immunity within that population.

4. Strategic Vaccination: Focused immunization of the at-risk cattle or areas when there is an epidemic.

Efforts to produce useful H5N1 vaccines that can be deployed for protection of the dairy cows are hampered by the high mutation rate of the virus, safety and efficacy testing, as wells as challenges involved in production and distribution of the vaccines (Okello, 2013).




Figure 4: Types of Influenza Vaccines6. Preventive Measures and Biosecurity

The awareness and control of H5N1 avian influenza from entering into and circulating in the dairy cows are essential to the health and productivity of the animals. This demands an integrated approach that accord much consideration to measures aimed at minimization of disease entry and proliferation within and between farms (Authority et al., 2018).

One of the major components of bio security is accommodation and layout of the farm. It was revealed that avoiding exposure of dairy cows to H5N1 sources such as wild birds and other poultry farms was effective in reducing transmission risks within the farm (Gao et al., 2024). This might entail construction of some barriers, housing of various animals separately, and placement of feed and water in appropriate areas.

Another crucial intervention is the necessity to control the access to the farm. Restricting only the management, workers, and vehicles importantly prevents the circulation of contaminated persons and items. It can also be supported by stricter measures enforced at the time of entry and exit into the premises, washing or disinfection of vehicles and footwear (Wireman, 2003).

Survival of H5N1 on equipment, vehicle, and facilities must be minimized through regular cleaning and disinfection. This is even more crucial especially where the input belongs to more than one farm or different personnel. It is important that the standards of cleaning and disinfection should be instituted and practiced so as to enhance the efficiency of these practices.

Measures of quarantine are significant as they help avoid the set in of H5N1 into a dairy population. New animals, animals brought back from shows or markets and those with clinical signs consistent with diseases should be isolated and observed for 7 days before being integrated to the main group. This makes it possible to identify and quarantine people who have possibly been infected with the virus to help curb the outbreak (Yadav et al., 2020).

Another interesting feature of biosecurity is wildlife management. Wild birds are considered

as natural hosts of H5N1, especially water birds. Exclusion of wild birds from the farm through use of barriers such as nets, or prevent their approach through using repellants will help minimize contact with the dairy cows. Moreover, feed and water availability should also be safeguarded to prevent contact with wild birds (Feare, 2010).

It is, therefore, very crucial to make sure that feed and water are safe. Proper storage of feed at a safe place and away from birds especially the wild ones that spread H5N1 virus helps avoid contamination (Kelly et al., 2008). The aspects such as water testing and treatment can be a way of eradicating the spread of the virus through water sources.

The other factor that should not be neglected is hygiene of people working on the farms and the tools they use. Treating and ensuring the proper utilization of personal protective gears like boots, coveralls, and gloves when dealing with the H5N1 infected cows or handling objects that may have come into contact with infected cows would help in curbing the spread of the virus from the affected area of the farm to other sections. Cleaning techniques include regular washing of shared equipment such as milking machines, milking tools and utensils after use or when used with infected animals to reduce cross contamination (Organization, 2006).

Determination of H5N1 at an early stage is a crucial step to enable prompt action and prevent its spread. Dairy cows should be checked frequently for clinical symptoms of diseases that could show early signs of an outbreak of the disease, including respiratory distress, low productivity, and weird behavior (Poovorawan et al., 2013). PCR and other diagnostic tests can help detect the presence of the virus in such occurrences and ensure that control measures are taken immediately.

High levels of public awareness are important when it comes to the practice of biosecurity in human societies. The business side of combatting H5N1 includes making sure that all farm personnel undergo and understand biosecurity procedures, disease identification, and emergency response measures. Interaction with colleagues in the field of veterinary medicine, industry associations, and governmental institutions helps to achieve synergy in the fight against diseases.

7. Economic and Social Impacts

Epidemics of H5N1 avian influenza in infected and neighboring dairy farms bring significant social and economic impacts to these farms and the overall dairy farming sector (Proença-Módena et al., 2007).

Economic Impacts

Direct Economic Impact:

• Loss of Milk Production: Milk production is influenced and this in turn influences the amount of money that a farm will generate (Somda et al., 2005).

• Increased Costs: Other costs include veterinary treatment, drugs and improved breeds securities which are costly.

• Culling and Losses: If the animals are infected or have been exposed to the virus, they will be slaughtered; this will cause a loss of money.

Indirect Economic Impact:

• Market Disruptions: Concerns about the purity of milk during epidemic contribute to the economy since it leads to changes in the prices of milk and hitches in selling milk products.

• Supply Chain Challenges: The processors and distributors face a challenge when it comes to attaining and maintaining the right volume of milk supply.

• Long-Term Financial Stability: Any outbreak of the disease puts a very huge cost to the

dairy farms, and particularly to small scale farming businesses that can be greatly affected.

Social Implications

• Livelihood Threat: Most of the affected communities rely on dairy farming as their source of income hence, economic loses are against their welfare.

• Community Stress: Measures aimed at controlling and preventing outbreaks may lead to heightened anxiety and stress in farmers and communities that would need friends and resources for mental health.

• Education and Awareness: Knowledge about the H5N1 virus and biosecurity measures is an important component of capacity-building for farmers fighting the disease.

Policy and Governance

• Biosecurity Regulations: Governments have to offer various bio-security features like observation, immunization, and reporting of every disease.

• Financial Support: Assistance in the form of guarantees, subsidies or insurance might help dairy farmers to overcome such an economic blow.

• International Collaboration: Hence, collaboration and cooperation at the international level is important for early detection, follow up action and to prevent virus circulation across the borders.

Future Challenges and Strategies

• Research and Innovation: A lot of money is needed for research if better vaccines, improved diagnostic tools, and ways to contain the disease are to be produced.

• Community Resilience: This would educate, train and provide support networks so that as communities, they would be ready as well as be in a position to handle the effects of the outbreak.

• Adaptive Management: It is thus important to increase the adaptive capacity when dealing with change such as viral mutagenesis in dairy farming systems.

8. Future Directions and Research Needs

Dairy industry towards H5N1 influenza is still under pressure and so further studies and more developments are necessary for better handling of diseases control and preparedness for adverse effects (McLeod, 2010).

Emerging Trends

1. Antigenic Evolution: This is alongside the fact that mutation of the H5N1 virus is not only a one-off occurrence but this has led to the appearance of new strains that affect different species, in this instance, dairy cows, in different ways (Scoones, 2010).

2. Globalization and Trade: Circulation of products and animals across international borders helps in virus spread across borders which makes international cooperation to be very essential (Cartín-Rojas, 2012).

3. One Health Approach: It is due to the social acceptance of the One Health concept in addressing diseases that are realized to have human-animal-environmental interface aspects like zoonotic H5N1 (Zhou & Tanner, 2022).

Research Gaps

1. Vaccine Efficacy: As such, more researches need to be made for purposes of assessing the existing H5N1 vaccines being administered to the dairy cows on various aspects including on

the duration of immunity offered and cross protection against other strains.

2. Transmission Dynamics: In the context of such transmission paths, the dynamics of H5N1 within dairy farms and between several farms have to be appreciated so as to come up with a control mechanism.

3. Host Immune Response: Further investigations will be conducted on survival of H5N1 in dairy cow and the objective is to identify immunological markers of protective immunity against avian influenza and facilitate the development of new vaccines.

4. Antiviral Strategies: More detail research on new antiviral drugs and therapy may assist in finding out other therapies applicable in treating the affected cows.

Innovative Approaches

1. Genomic Surveillance: Such awareness of the genetic changes in the virus makes it easy to predict and also develop appropriate vaccines (Gardy & Loman, 2018).

2. Precision Vaccinology: Some suggestions of course, could be improved by stating the particular H5N1 strains for which the vaccine should be developed.

3. Digital Health Solutions: Increased mHealth utilization for reporting diseases and insights collection can improve surveillance and reaction.

Collaborative Initiatives

1. Public-Private Partnerships: COD Promotes: increasing collaboration with federal/state universities, industry and others in the areas of communication, research & grant support, funding and technology transfer.

2. International Cooperation: The studies have it that the schemes which foster dissemination of information, development of capacity among the partners, and collaboration in research enhance the global level of preparedness.

3. Community Engagement: Efforts can therefore be made to ensure that farmers and other members of the communities participate in research and education so as to take up the task of combating diseases.

Future Directions

1. Enhanced Surveillance: Encouraging the application of various surveillance systems to pinpoint and alert the disease and likely occurrences early enough.

2. Capacity Building: The preparedness of diagnostic laboratories, veterinary services, and public health structures that have to be strengthened for increasing the capacities of response.

3. Resilience Building: Increase support from extension agents; educate; also devise contingencies for the dairy farming and financial scenarios in case of an outbreak.

9. Summary

The threat of H5N1 influenza is even higher for dairy cows, as the problem is quite diverse and requires not only local but also systemic solutions to prevent and deal with. They concluded that the virus presents severe economic risks to dairy farming such that milk production lowers, operation costs raise, and the market situation becomes fragile. The disease also has far reaching consequences economic, both for farmers and welfare of rural folks. Vaccination, identification of infected birds promptly, and other matters that pertain bio-security measures are essential to reduce the transmission of H5N1 and hence the impact in the different dairy herds.

Improving global networking and research funding are vital to the creation of better vaccines, diagnostic measures, and ways to address viral diseases. Genomic surveillance precision vaccinology and digital health solutions point to potentially major directions to enhance disease prevention and control. Moreover, preparing the dairy farming communities for disasters and building their capacity to cope with adversity is also an important proactive step. According to the principle of One Health and development of cooperation with the stakeholders it may be speculated that the effectiveness of H5N1 outbreak preventing, detecting and responding would be enhanced and animal health, food security as well as public health globally may be preserved.

References

Abdelwhab, E., & Hafez, H. M. (2012). Insight into alternative approaches for control of avian influenza in poultry, with emphasis on highly pathogenic H5N1. Viruses, 4(11), 3179-3208.

Alqazlan, N., Astill, J., Raj, S., & Sharif, S. (2022). Strategies for enhancing immunity against avian influenza virus in chickens: A review. Avian Pathology, 51(3), 211-235.

Authority, E. F. S., Prevention, E. C. f. D., Control, Influenza, E. U. R. L. f. A., Adlhoch, C., Brouwer, A., Kuiken, T., Mulatti, P., Smietanka, K., Staubach, C., & Muñoz Guajardo, I. (2018). Avian influenza overview February–May 2018. Efsa Journal, 16(6), e05358.

Bauzile, B., Sicard, G., Guinat, C., Andraud, M., Rose, N., Hammami, P., Durand, B., Paul, M. C., & Vergne, T. (2022). Unravelling direct and indirect contact patterns between duck farms in France and their association with the 2016–2017 epidemic of Highly Pathogenic Avian Influenza (H5N8). Preventive Veterinary Medicine, 198, 105548.

Beerens, N., Germeraad, E. A., Venema, S., Verheij, E., Pritz-Verschuren, S. B., & Gonzales, J. L. (2021). Comparative pathogenicity and environmental transmission of recent highly pathogenic avian influenza H5 viruses. Emerging microbes & infections, 10(1), 97-108.

Bhalerao, U., Mavi, A. K., Manglic, S., Sakshi, Chowdhury, S., Kumar, U., & Rohil, V. (2023). An Updated Review on Influenza Viruses. Emerging Human Viral Diseases, Volume I: Respiratory and Haemorrhagic Fever, 71-106.

Cartín-Rojas, A. (2012). Transboundary animal diseases and international trade. International Trade from Economic and Policy Perspective, 143, 166.

Duarte, P., El-Nakeep, S., Shayestegan, F., Tazerji, S., Malik, Y., Roncada, P., Tilocca, B., Gharieb, R., Hogan, U., & Ahmadi, H. (2024). Addressing the recent transmission of H5N1 to new animal species and humans, warning of the risks and its relevance in One-Health. Ger. J. Microbiol, 4(2), 39-53.

Feare, C. J. (2010). Role of wild birds in the spread of highly pathogenic avian influenza virus H5N1 and implications for global surveillance. Avian diseases, 54(s1), 201-212.

Gao, R., Sreenivasan, C. C., Feng, C., Li, F., & Wang, D. (2024). Cross-Species Transmission of Highly Pathogenic Avian Influenza (HPAI) H5N1 Virus in the US Dairy Cattle: A Comprehensive Review.

Gardy, J. L., & Loman, N. J. (2018). Towards a genomics-informed, real-time, global pathogen surveillance system. Nature Reviews Genetics, 19(1), 9-20.

Gisbert, J., Menchén, L., García-Sánchez, V., Marín, I., Villagrasa, J., & Chaparro, M. (2012). Comparison of the effectiveness of two protocols for vaccination (standard and double dosage) against hepatitis B virus in patients with inflammatory bowel disease. Alimentary pharmacology & therapeutics, 35(12), 1379-1385.

Graziosi, G., Lupini, C., Catelli, E., & Carnaccini, S. (2024). Highly pathogenic avian influenza (HPAI) H5 clade 2.3. 4.4 b virus infection in birds and mammals. Animals, 14(9), 1372.

Gulati, A., & Juneja, R. (2023). Poultry Revolution in India: Lessons for smallholder production systems.

Huang, P., Sun, L., Li, J., Wu, Q., Rezaei, N., Jiang, S., & Pan, C. (2023). Potential cross-species transmission of highly pathogenic avian influenza H5 subtype (HPAI H5) viruses to humans calls for the development of H5-specific and universal influenza vaccines. Cell discovery, 9(1), 58.

Kappes, A., Tozooneyi, T., Shakil, G., Railey, A. F., McIntyre, K. M., Mayberry, D. E., Rushton, J., Pendell, D. L., & Marsh, T. L. (2023). Livestock health and disease economics: a scoping review of selected literature. Frontiers in veterinary science, 10, 1168649.

Kelly, T. R., Hawkins, M. G., Sandrock, C. E., & Boyce, W. M. (2008). A review of highly pathogenic avian influenza in birds, with an emphasis on Asian H5N1 and recommendations for prevention and control. Journal of avian medicine and surgery, 22(1), 1-16.

Kim, H.-R., Park, C.-K., Lee, Y.-J., Woo, G.-H., Lee, K.-K., Oem, J.-K., Kim, S.-H., Jean, Y.-H., Bae, Y.-C., & Yoon, S.-S. (2010). An outbreak of highly pathogenic H5N1 avian influenza in Korea, 2008. Veterinary microbiology, 141(3-4), 362-366.

Lee, H. Y., Topham, D. J., Park, S. Y., Hollenbaugh, J., Treanor, J., Mosmann, T. R., Jin, X., Ward, B. M., Miao, H., & Holden-Wiltse, J. (2009). Simulation and prediction of the adaptive immune response to influenza A virus infection. Journal of virology, 83(14), 7151-7165.

Ly, H. (2024). Highly pathogenic avian influenza H5N1 virus infections of dairy cattle and livestock handlers in the United States of America. In (Vol. 15, pp. 2343931): Taylor & Francis.

Marrana, M. (2022). Epidemiology of disease through the interactions between humans, domestic animals, and wildlife. In One Health (pp. 73-111). Elsevier.

McLeod, A. (2010). Economics of avian influenza management and control in a world with competing agendas. Avian diseases, 54(s1), 374-379.

Okello, A. L. (2013). Beyond avian influenza: policy considerations for the implementation of a “one health” approach in developing countries.

Organization, W. H. (2006). Collecting, preserving and shipping specimens for the diagnosis of avian influenza A (H5N1) virus infection: guide for field operations.

Peiris, J. S. M., Cheung, C. Y., Leung, C. Y. H., & Nicholls, J. M. (2009). Innate immune responses to influenza A H5N1: friend or foe? Trends in immunology, 30(12), 574-584.

Polley, S., Biswas, S., Kesh, S. S., Maity, A., & Batabyal, S. (2022). The link between animal manure and zoonotic disease. In Animal Manure: Agricultural and Biotechnological Applications (pp. 297-333). Springer.

Poovorawan, Y., Pyungporn, S., Prachayangprecha, S., & Makkoch, J. (2013). Global alert to avian influenza virus infection: from H5N1 to H7N9. Pathogens and global health, 107(5), 217-223.

Proença-Módena, J. L., Macedo, I. S., & Arruda, E. (2007). H5N1 avian influenza virus: an overview. Brazilian Journal of Infectious Diseases, 11, 125-133.

Rosone, F., Bonfante, F., Sala, M. G., Maniero, S., Cersini, A., Ricci, I., Garofalo, L., Caciolo, D., Denisi, A., & Napolitan, A. (2023). Seroconversion of a swine herd in a free-range rural multi-species farm against HPAI H5N1 2.3. 4.4 b clade virus. Microorganisms, 11(5), 1162.

Schat, K. A., & Skinner, M. A. (2022). Avian immunosuppressive diseases and immune

evasion. In Avian immunology (pp. 387-417). Elsevier.

Scoones, I. (2010). Avian influenza: science, policy and politics. Routledge.

Shi, J., Zeng, X., Cui, P., Yan, C., & Chen, H. (2023). Alarming situation of emerging H5 and H7 avian influenza and effective control strategies. Emerging microbes & infections, 12(1), 2155072.

Smith, O. M., Snyder, W. E., & Owen, J. P. (2020). Are we overestimating risk of enteric pathogen spillover from wild birds to humans? Biological Reviews, 95(3), 652-679.

Somda, J., Kamuanga, M., & Tollens, E. (2005). Characteristics and economic viability of milk production in the smallholder farming systems in The Gambia. Agricultural systems, 85(1), 42-58.

Trevisi, E., & Minuti, A. (2018). Assessment of the innate immune response in the periparturient cow. Research in veterinary science, 116, 47-54.

Velkov, T., Ong, C., Baker, M. A., Kim, H., Li, J., Nation, R. L., Huang, J. X., Cooper, M. A., & Rockman, S. (2013). The antigenic architecture of the hemagglutinin of influenza H5N1 viruses. Molecular immunology, 56(4), 705-719.

Wireman, T. (2003). Maintenance management and regulatory compliance strategies. Industrial Press Inc.

Yadav, M. P., Singh, R. K., & Malik, Y. S. (2020). Emerging and transboundary animal viral diseases: Perspectives and preparedness. Emerging and transboundary animal viruses, 1-25.

Yoo, D.-S., Lee, K.-N., Chun, B.-C., Lee, H.-S., Park, H., & Kim, J.-K. (2022). Preventive effect of on-farm biosecurity practices against highly pathogenic avian influenza (HPAI) H5N6 infection on commercial layer farms in the Republic of Korea during the 2016-17 epidemic: A case-control study. Preventive Veterinary Medicine, 199, 105556.

Zappulli, V., Ferro, S., Bonsembiante, F., Brocca, G., Calore, A., Cavicchioli, L., Centelleghe, C., Corazzola, G., De Vreese, S., & Gelain, M. E. (2020). Pathology of coronavirus infections: A review of lesions in animals in the one-health perspective. Animals, 10(12), 2377.

Zhou, X.-N., & Tanner, M. (2022). Science in one health: a new journal with a new approach. In (Vol. 1, pp. 100001): Elsevier.

About the Authors

Arifa Mehreen is an Assistant Professor of Zoology at the University of Agriculture Faisalabad (UAF). Her research focuses on antimicrobial properties of natural products, drug formulation and pharmacokinetics, and characterization of bacterial pathogens. She has studied the antimicrobial and toxicological effects of Origanum vulgare, developed a fixed-dose combination drug for cardiovascular disease, and investigated the characteristics of Staphylococcus aureus strains.

E-mail arifa.mehreen@uaf.edu.pk ORCID 0000-0001-5899-8027

Muhammad Majeebullah holds an M.Phil. degree in Zoology from the University of Agriculture Faisalabad (UAF), Pakistan. His research focuses on the effects of nanoparticles on growth, contributing to advancements in this emerging field. Muhammad is dedicated to exploring how nanoparticles can influence biological processes, aiming to uncover new insights and applications.

Email muhammadmajeebullah336@gmail.com ORCID0009-0002-2138-2039

Shagufta Abbas is a Ph.D. scholar at The Women University Multan, Pakistan. Her research is centered on neuroscience, with a keen interest in understanding the complexities of the brain and nervous system. Shagufta is committed to advancing knowledge in this field through her innovative research. Her work aims to contribute significantly to the scientific community.

Email shaguftaabbas6373@gmail.com ORCID 0000-0002-8161-3671

Dr. Muhammad SAFDAR earned his PhD in molecular biology and genetics from Gaziantep University, Turkey. He is a faculty member in the Breeding and Genetics department at Cholistan University of Veterinary and Animal Sciences (CUVAS), Bahawalpur, Pakistan. His research interests are molecular genetics and genomics, nutrigenomics, nano-genomics, bioinformatics, biotechnology, and their applications. He has published more than 70 research articles in national and international journals. He has also written many book chapters as well as an edited book. He is an associate editor for international journals.

E-mail msafdar@cuvas.edu.pk , ORCID 0000 0002 3720 2090

Dr. Muhammad SAFDAR earned his PhD in molecular biology and genetics from Gaziantep University, Turkey. He is a lecturer in the Breeding and Genetics department at Cholistan University of Veterinary and Animal Sciences (CUVAS), Bahawalpur, Pakistan. His research interests are molecular genetics and genomics, nutrigenomics, nano-genomics, bioinformatics, biotechnology, and their applications. He has published more than 70 research articles in national and international journals. He has also written many book chapters as well as an edited book. He is an associate editor for international journals.

E-mail msafdar@cuvas.edu.pk , ORCID 0000 0002 3720 2090

To Cite This Chapter

Mehreen, A, et al. (2024). IMPACT OF H5N1 INFLUENZA ON DAIRY COWS. In Animal Production and Health (pp.320-332). ISRES Publishing.