

Maham Shahzadi is a final-year BS Biotechnology student at Cholistan University of Veterinary and Animal Sciences, Bahawalpur (CUVAS). She is passionate about exploring the diverse applications of biotechnology to improve human health, agriculture, and marine ecosystems. Her research interests include developing novel diagnostics and therapeutics, enhancing crop yields and sustainability, and utilizing marine resources for drug discovery and environmental protection. With a strong academic background and a keen interest in cutting-edge research, Maham is poised to make significant contributions to the field of biotechnology.

E-mail: mahamshahzadi005@gmail.com

ORCID: 0009-0005-9868-2162

Shafeeq Ur Rehman earned his Bachelor in Applied microbiology from Cholistan University of Veterinary and Animal Sciences (CUVAS) Bahawalpur Pakistan. Currently he is pursuing M. phil in Microbiology from Cholistan University of Veterinary and Animal Sciences (CUVAS) Bahawalpur Pakistan. He is also working as research assistant in a project titled as “Robust Molecular Detection of Silent Circulation of FMD and PPR Viruses in Cholistan as a Model for Disease-Free Zone”. His research interest is in FMD and PPR Viruses, Mucosal Immunity, Virology, Molecular cell biology, Cell culture, Microbial genetics and CRISPR case tools for combating viral diseases. He has published 7 articles and many book chapters in national and international journals.

Email: shafeequrrehmanbhutta@gmail.com

ORCID: 0000-0003-3571-8226

Muhammad Ejaz earned Bachelor's in applied microbiology from Cholistan university of veterinary and animal sciences (CUVAS) Bahawalpur Pakistan. His research interest in Microbial genetics, Molecular biology, and food quality and its application.

Email: ijazrasheed334@gmail.com

ORCID: 0009-0006-1709-6653

Dr. Muhammad SAFDAR earned his PhD in Molecular Biology and Genetics from Gaziantep University, Turkey. He is Lecturer in the Breeding and Genetics department at Cholistan University of Veterinary and Animal Sciences (CUVAS), Bahawalpur, Pakistan. His research interests are molecular genetics and genomics, nutrigenomics, nano-genomics, bioinformatics, biotechnology, and their applications. He has published more than 70 research articles in national and international journals. He has also written many book chapters as well as an edited book. He is an associate editor for international journals.

E-mail: msafdar@cuvas.edu.pk

ORCID: 0000 0002 3720 2090

To Cite This Chapter

NAZIR M et al., (2024). Breaking barriers: advancements in genomic research and crispr technology. In *Current Studies in Health and Life Sciences 2024* (pp.196-210). ISRES Publishing.

Stem Cell Research: Unlocking The Potential for Regenerative Therapies

Maria NAZIR

Hameer Khan KHASKHELI

Ammara HAMEED

Tasawar ABBAS

Danish RIAZ

Shafeeq Ur REHMAN

Muhammad SAFDAR

1. Introduction

Stem cells have emerged as one of the brightest stars on the map, directing the course of modern medicine, and have attracted the attention of scientists as well as the common people. Such strategies of using the natural self-healing efficiencies of body tissues for curing such ailments is a new chapter in the history of medical science; it can be inferred that various diseases and injuries can be cured using the remedial advantages of regenerative medicine (Sorensen et al., 2022). In addition to this field, stem cells are some of the most exciting cells in biology owing to their ability to continue dividing for a rather long time and differentiate into many specialized cells (Hall & Watt, 1989). This double-faced characteristic of stem cells makes them useful resources for the treatment and regeneration of tissues, hence offering hope to patients with diseases that previously could not be treated. For instance, for neurodegenerative diseases such as Parkinson's and Alzheimer's disease, cardiovascular diseases, heart failure, and myocardial infarction stem cell therapy can be used in almost all the diseases that humans are unconsciously susceptible (Dawson et al., 1995). Thus, once people understand how different elements of stem cells' environment influence these cells, the scientific community will be paving the way for producing therapeutic approaches that could help millions of people with chronic pains and heal diseases as well as prolong human lifespan.

Scientists discovered that cell lineages could differentiate into all blood cells via the bone marrow hematopoietic stem cells (HSCs). It formulated the guidelines for bone marrow transplantation the only treatment for some forms of blood cancer and other hematological disorders. Stem cells were regarded as a priceless biomaterial; however, these cells were analyzed and researched in only several variations up to the end of the period of the nineteen nineties when hESCs were derived and cultured (Ericson, 2007). ESCs arise from the blastocyst and are noted for their Pluripotency or the ability of these cells to differentiate into any cell of the body, a process they derive from the blastocyst. This discovery gave way to subsequent surveys because people understood that stem cells are effective in solving different human diseases owing to their ability to repair tissues and organs. As a result of improving technological advancement and enhanced knowledge of stem cells, stem cell research has been boosted maximally over the last decade. However, with the later development of induced pluripotent stem cells (iPSCs), which is made by transforming an ordinary body cell into that which is almost embryonic, ethical considerations (Lewandowski & Kurpisz, 2016) that come with the use of ESCs are eliminated, although their totipotency is not opposed. It has allowed doctors to prescribe medication dependent on the personal traits of the patient, allow researchers to construct models of diseases, along with finding new medications. According to Shi et al., the consequent patient-specific nature of iPS cells enables the researcher to create a disease model in a dish, investigate the disease progression, and trial the drugs in a far more realistic setting. Besides, they serve purposes in regenerative medicine because the cells can be terminal to other types of cells for transplantations and tissue replacement (Ringe et al., 2002).

Moreover, findings have been made and reported that provide information on identifying numerous subpopulations of adult stem cells in different body parts. They contain tissue-specified stem cells that play vital roles in the homeostasis and repair of tissues, and the element potential of these tissues is discussed based on several diseases (Beerman & Rossi, 2015). For instance, MSCs derived from bone marrow have therapeutic use in autoimmune-related disorders, new blood vessel formation, and the replacement of damaged vital organ tissues. NSCs present the capability to replace flawed neurons and improve rehabilitation in global ischemia, neurological disorders, and spinal cord injury (Vishwakarma et al., 2014). Stem cell research has not been in vain as it extends direct medical application. At present, hematopoietic stem cell transplantation is known as a therapy for multiple blood diseases and kinds of cancer, and MSCs derived from bone marrow are used in autoimmune diseases and tissue regeneration. In ophthalmology, ESCs have been applied to produce layers of RPE in the treatment of one of the most prevalent causes of blindness, AMD (Schwartz et al., 2016). In this chapter, the reader will be first acquainted with the subject of stem cell research, especially emphasizing the differentiation of stem cells; they will also get a preview of the other types of stem cells, and, finally, the biological activities of stem cells will be explained in detail. Self-renewal and multipotent differentiation potential cells are known as stem cells; the different techniques used for isolation culture and manipulations of the stem cells and the development and advances made in the implementation of ideas into operation shall be understood. Thus, in the context of stem cells, let us clarify the position as to how we when it comes to confronting the current difficulties, are to shed light on the road that will incorporate regenerative medicine into a truly global picture beneficial for millions of people on earth.

2. Types of Stem Cells

Depending on their origin and differentiation potential, stem cells are categorically recognized in many categories. Since cells of different types have unique features and roles, it is crucial to understand them from the perspective of using stem cells in the literature of regenerative medicine.

2. 1 The first one develops from the Blastocyst

ESCs are isolated from the inner cell mass of early-stage embryos, preferably at the blastocyst stage, which is between five to seven days after fertilization (Desmarais et al., 2011). These cells are pluripotent, meaning they have the remarkable ability to differentiate into any cell type in the adult body, giving rise to all three germ layers. Some of the interesting features of stem cells include the ectoderm, mesoderm, and endoderm layer in humans. This makes ESCs versatile and this attribute is considered a plus in regenerative medicine since it can replace tissues that are destroyed or ravaged by diseases in the various body organs and systems (Figure 1). For instance, the scientific technique has been tested, and therefore, it is possible to differentiate ESCs into insulin-producing pancreatic beta cells to combat type 1 diabetes (Soria et al., 2001). However, the use of ESCs has been controversial because of its source, which is human embryos, a process most people consider to be eradicating human life. Even if an effort has been made to derive ESCs from other sources, such as somatic cell nuclear transfer (SCNT), the methods are still experimental and thus not clinically practicable (Šarić et al., 2010).




Figure 1. Human Embryonic Cell Differentiation

2. 2 Adult Stem Cells or Mesenchymal Stem Cells

Stem Cells These are stem cells that are taken from adults and are also known as postnatal stem cells are found in different parts of the body. This includes the bone marrow, adipose tissue of an adult, and umbilical cord blood of a newborn baby (Malgieri et al., 2010). Like any other stem cell, therefore, ESCs are pluripotent in their capabilities of differentiating into the various major cell types of an organism, while ASCs are multipotent, as many of the researchers would like to believe, by having the capability of differentiating into not more than three or four cell types of their lineage. For example, hematopoietic stem cells are found in the bone marrow, which can give rise to all the blood cells, or the mesenchymal stem which can transform into a bone cartilage or adipose tissue cell (Figure 2).




Figure 2. Adult Stem Cell differentiates to become Mesenchymal Stem Cell.

However, unlike ESCs, ASCs have certain advantages because of their low differentiation potential (Uzbas et al., 2015). They can be derived from adult people without ethical problems and can be reimplanted back into the body without facing rejection from the human body, unlike the ESCs. Moreover, some of the research has come across that some of these ASCs may have a specific property called the plasticity capacity through which it develops other cellular structures different from its source tissue (Sordi et al., 2017).

2. 3 Human embryonic stem cells (hESCs)

This involved using some chemicals to get a somatic cell to aspire to become an iPSC during the body's development processes. These cells are derived from normal somatic cells such as skin or blood cells and reprogrammed back to this state of embryonic stem cells through the agency of four genes, namely Oct4, Sox2, Klf4, and c-Myc (Zakrzewski et al., 2019). iPSCs are also known as pluripotent stem cells, just like ESCs, and this simply means that they can turn into any type of cell in the body as and when required. However, iPSCs do not come with some of the concerns of ESCs, for they can be differentiated from the easily obtainable adult cells. In addition, one can derive iPSCs from patients for disease studies, drug testing, and maybe possible regenerative medicine for each single patient. For example, researchers have used iPSCs of affected patients with genetic diseases; they observed the advancement of the disease on a plate and were able to establish possible therapeutic interventions (Bonaventura et al., 2021).

Nevertheless, the reprogramming process is related to altering genetic and epigenetic statuses, which affects the safety and function of iPSCs in clinical use. More research has to be done, and thereby, the quality control of the studies to establish how such matters can be handled to allow the

safe introduction of iPSC-based therapies into practice (Jha et al., 2021). Thus, all types of stem cells are different from one another by their peculiarities and treatments, which scientists have opened are almost infinite. Analyzing how the two human body parts have their individuality and how science is attempting to use stem cells and their capacity to self-renew, scientists are now able to come up with tonics that could go a long way in satisfying many medical needs.

3. Stem Cell Biology and Mechanisms

Stem cells are a particular kind of cell that possess different nature and characteristics from other cells. This paper describes the principal regulators of stem cell biology, which is fundamental information in stem cell-based therapeutic initiatives.

3. 1 Stem Cell Niches

Stem cells reside in shelters known as niches that are fundamental in the regulation of stem cells' properties. These niches provide mechanical substrate and also various bioactive molecules, cytokines, and factors of the extracellular matrix that control stem cell functions such as self-renewal, differentiation, and quiescence (Singh et al., 2019). Stem cell niche structure and organization can also be rather diverse concerning stem cells' overall tissue and differentiation. For example, the precise locality of HSCs is the bone marrow containing osteoblasts, endothelial cells, perivascular cells, and sympathetic neurons (Birbrair & Frenette, 2016). Known as niche cells, these cells employ various signals mostly through Notch/Wnt, SDF-1/CXCR4, and angiopoietin-1/Tie2 to regulate HSCs self-renewal, differentiation, and release (Portale, 2018). They are situated at the bottom of the crypts of Lieberkuhn and consist of Paneth cells and Lgr5 + stem cells (Schewe & Fodde, 2018). Despite wnt signaling for the renewing of the intestinal stem cells, notch signaling is not involved in the transmutation into various lineages of cells in the intestine (Foster et al., 2002). The stem cell niche is found to function as the master circuit, keeping a delicate balance between the stem cells' infinite division and differentiation. It improves the initiation of stem cell availability in the process of replacing tissues as individual life and also serves as a map for stem cells in case they are required to transform into certain cells for the formation of tissues, more so, for replacement and repair purposes (Saltzman, 2004). The changes of microenvironmental stimuli in the micro-territory fragment can lead to deviations in stem cell behavior and their participation in the diseases' progress, like cancers and degenerative diseases. For instance, the study highlighted how the modification of the bone marrow's place of dwelling leads to leukemia as well as other types of hematology-borne cancer.

3. 2 Microenvironments and Signaling Pathways

The stem cell niche is an active area summarized as the amalgamation of several signals that affect the stem cell functionality. These signaling pathways (Figure 3) include autocrine where the signals are synthesized and released by stem cells or paracrine factors coming from other stem cells or the surrounding matrix (Burchfield & Dimmeler, 2008). Some of the key signaling pathways involved in stem cell regulation include:




Figure 3. Different Signal Transduction Pathways

• **Notch signaling:** This pathway plays an especially critical role in the decisions that are made about cell lineage and is beneficial in maintaining the stem cells being ‘pluripotent’ while at the same time promoting differentiation into certain cells/tissues (Walia et al., 2012). For the hematopoietic system, Notch looked into the fate map of HSC concerning either maintaining cell proliferation or differentiating into the lymphoid or the myeloid lineage.

• **Wnt signaling:** The Wnt signaling pathway is involved in the regular development of an embryo and for the maintenance of tissue homeostasis and is essential for stem cell self-renewal, proliferation, and differentiation in different body tissues (Sokol, 2011). The Wnt signaling pathway helps in maintaining the ES and floor of the mouth iPS cells ' self-renewable states, while the inhibition of the Wnt signaling pathway plays a significant role in the differentiation process.

• **BMP signaling:** This has been relayed in diverse cell actions, including cell division, stem cell differentiation, and cell death, and is core to stem cell differentiation in various tissues. In the context of the bone marrow niche, BMP signaling assists in the same by restricting the HSCs into quiescence and hence assists in rewinding the long-term repopulating potential of these cells (Burel, 2022).

• **Hedgehog signaling:** In addition, this pathway participates in embryonic development and tissue stimulation and regulates the stem cell reaction in various tissues. In the skin, hedgehog signaling functions to retain Hair Follicle Stem cells in the G0 phase and regulate the emergence of hair structures (Zhang & Chen, 2024).

• **FGF signaling:** This pathway is used in processes like growth, development of cells, and movement and regulates stem cell activity in specific body regions. Anything that affects balanced FGF signaling directly affects ESCs and iPSCs’ capacity to maintain their original embryonic stem cell state and their capacity to differentiate into lineages of the human body (Mong, 2013).

These signaling pathways can be interconnected, and this is how one obtains vast signal transduction networks that control stem cell functions in an idiosyncratic cell type and developmental phase. It is, therefore, important to understand these multiple interactions so that more precise treatments can be developed to modulate stem cell activity for therapeutic purposes.

3. 3 Differentiation and Plasticity

Differentiation is capable of specializing and supercilious the functions of the cells of the tissues and organelles. This process of differentiation is hereby described as a well-coordinated series of molecular changes in the cell, the pattern of genes, protein synthesis, and cell structure.

Stem cell's ability to respond to signals that cause their differentiation

This process of stem cell differentiation is a seemingly complex process that demarcates the genes to be turned on and the genes to be turned off through several factors, including both the intrinsic and the extrinsic factors. Modifications such as DNA methylation and histone acetylation alterations are used in the regulation of the gene in differentiation (Ikegami et al., 2009). Among the local key players, the most essential consists of those proteins capable of binding DNA sequences, which are, as a result, termed transcription factors because of their ability to switch target genes' activation or repression (Suter, 2020). Stem cells usually act according to the differentiation hierarchy; therefore, higher-order multipotent stem cells can differentiate in several cell types of a given lineage. In the process of differentiation, stem cell acquires partial characteristics of differentiation, which are prep producer cells for a particular lineage, but can go further to develop still other, more specialized IP cells before getting completely differentiated. For example, HSC in the hematopoietic system can differentiate to PCM or CLP, which in turn differentiate to virtually all blood cells such as red blood cells, the plates, as well as immune systems (Ranjitha, 2019). This implies that doing the differentiation process does not necessarily warrant this aspect of differentiation. Some stem cells are fully plastic because they either de-differentiate or trans-differentiate into other cell types under certain conditions. For instance, MSCs have been reported to trans-differentiate into ANLL-like cells, a discovery that should enable the application of such cells in the management of neurological diseases (Karpowicz, 2004). However, as regards the future of stem cells and their capability, this is still exceptional.

Factors Influencing Lineage Specification

The decision of which pathway to take is an action carried out by the stem cell through intrinsic and extrinsic processes. Other concurrent exogenous aspects include the epigenetic state of the stem cell at the onset and the switching on certain transcriptional factors as a prognosis of the stem cell's destination. The surrounding microenvironment, such as neighboring cells, signaling molecules, growth factors, and parts of the extracellular matrix, are the ones that provide the necessary signal to make destiny commitments (Lloyd-Lewis et al., 2019). The relative intrinsic and extrinsic force rates are pegged on certain environmental stimuli factors such as oxygen partial pressure, nutrient delivery, and biomechanical application. These can elicit so many responses in a cell that will spur the stem cell to either encourage or suppress the genes within and consequently determine its fate. For instance, hypoxia has been known to promote changes of MSCs to osteoblasts with the activation of nuclear factor HIF-1 (Ejtehadifar et al., 2015). Much effort has hence been directed at evaluating the complex signal transduction processes that guide stem cell differentiation as a way of rightly controlling the processes in a bid to provide the much-needed directions for innovative stem cell-based therapies.

3. 4 Self-Renewal and Proliferation

Stem cells can be released through self-renewal. This is the process of differentiating more stem cells that are phenotypically close to the original stem cell (Zipori, 2005). It is an important process of how the stock of stem cells in an individual is regenerated during a lifetime to enable tissue self-repair. However, at the same time, the uncontrolled capacity for self-renewal may lead to the formation of tumors and other diseases. This process of self-renewal and differentiation is controlled in a very similar way and helps to balance the tissue without the elimination of stem cells or an excessive amount of cell amplification. This balance is achieved via the presence of signaling pathways in the cell and additionally through transcription factors and epigenetic modifiers.

Molecular Pathways Regulating Self-Renewal

The records of the molecules, which include signals involved in the regulation of stem cell self-renewal, include the following. The Wnt signaling pathway is also used to control self-renewal capability, mainly in ESCs and ASCs (Jayaraman et al., 2021). Wnt signaling is useful in preserving the pluripotency of the cells due to the later activation of target genes such as c-Myc and cyclin D1, which are known genes that hinder the differentiation process (Sidrat et al., 2021). Furthermore, there are integrations of reprogramming pathways like Wnt with other signaling pathways such as Notch and TGF- β are involved in the regulation of the self-renewal process (Gordeeva, 2019). Another pathway that is also involved in stem cell self-renewal is the notched signaling pathway. While Hes1 belongs to the differentiation genes, it is, on the contrary, inhibited, while Bmi1, which is associated with stem cell self-renewal, is enhanced by NOTCH. This pathway is employed for the generation of renewal of HSCs, NSCs, and ISCs (Al-Jedai, 2008). Specifically well-described pathway that helps in stem cell survival, proliferation, and self-renewal is the PI3K/Akt signaling that is triggered by growth factors or cytokine. Downstream targets are launched, including the kinase mTOR (mechanistic target of rapamycin) mapped to photosynthesis and cell multiplication, besides FOXO transcription factors concerning genes associated with cell division and stress (Verma & Chatterjee, 2009). The same can be said about the ratio between the number of stem cells, in this case of the HaCaT cell line, and the differentiation of the cells at hand (Çetin et al., 2023). That is why regulating the stem cell cycle and determining the cell's developmental plan is critical for the tissue to remain stable. Therefore, cells of the body produce what might be referred to as an alarm in the form of inflammation whenever there is bodily tissue injury or damage, and the stem cells reproduce and begin to differentiate into a specific nutrient in a bid to replenish damaged lost cells. However, when these cells divide overly, they can then result in the formation of tumors or, if the division rate is inadequate, this could lead to tissue degeneration (Baba & Câtoi, 2007).

Several positive feedbacks avoid too numerous copies of the same cell type and negative feedbacks define the final fate of each given cell. Among these growth regulatory proteins, cyclins and cyclin-dependent kinases, and p 53 and Rb, the transcription factors that set the rate of cell division and the differentiation genes. HEPA growth factors such as EGF and FGF; cytokines such as IL-6, and cell adhesion molecules to get into the cell cycle and differentiate or to remain stem cells (Rhind & Russell, 2012). Hence, it bears a pertinent role as the stem cell niche in terms of right imposing stem cells' self-renewal and differentiation conditions. The niche signals may advise stem cell quiescence, which is a process of rather temporary withdrawal from the cell cycle with the intention of not depleting stem cells. Several processes attain this; creating inhibitory cytokines, including TGF- β , and obtaining quiescence-inducing niche cells (Moses, 2016). If tissue repair is needed, then the niche signals help the stem cells to proliferate and differentiate to replace the damaged or dead cells.

4. Methods Used in Stem Cell Research

This is because various procedures have been created for stem cell research. Through these procedures, stem cells can be isolated, grown, and manipulated. These techniques have assisted in enhancing knowledge of stem cell biology and given the basis for potential therapeutic applications.

Table 1: A list of methods that are employed in stem cells for isolation and culturing (Khan et al., 2018).

Technique	Stem Cell Type	Isolation Method	Culturing Method	Advantages	Limitations
Embryo Dissection	Embryonic Stem Cells (ESCs)	Manual dissection, laser dissection, immunosurgery	Feeder-dependent or feeder-free systems on specialized media	High pluripotency; Unlimited self-renewal potential	Ethical concerns; Risk of teratoma formation; Potential for immune rejection
Bone Marrow Aspiration/Biopsy	Hematopoietic Stem Cells (HSCs)	Aspiration from iliac crest or sternum; Biopsy from iliac crest	Liquid cultures with growth factors; Co-culture with stromal cells	Relatively easy to obtain; Well-established protocols; No ethical concerns	Limited differentiation potential (blood cells only); Invasive procedure
Mobilized Peripheral Blood Collection	HSCs	G-CSF administration followed by apheresis (blood separation)	Similar to bone marrow culture	Less invasive than bone marrow aspiration; Higher yield of HSCs	Requires G-CSF administration; Possible side effects
Adipose Tissue Harvest	Mesenchymal Stem Cells (MSCs)	Liposuction or surgical excision	Adherent cultures on specialized media	Abundant source; Easy to obtain; Minimal ethical concerns	Lower differentiation potential compared to ESCs; Potential for contamination
Cell Reprogramming	Induced Pluripotent Stem Cells (iPSCs)	Transfection or transduction of reprogramming factors (Oct4, Sox2, Klf4, c-Myc) into somatic cells	Similar to ESC culture	Avoids ethical concerns of ESCs; Patient-specific cells possible; No immune rejection	Reprogramming efficiency can be low; Risk of genetic abnormalities; Concerns about long-term safety
Umbilical Cord Blood Banking	HSCs, MSCs	Collection of cord blood after birth	Cryopreservation for future use	Rich source of stem cells; Non-invasive; Easily accessible; No ethical concerns	Limited quantity available; Not suitable for all patients

4. 1. Biopsy and Enrichment of Stem Cell Samples

Another procedure that is usually traditional but fundamental in stem cell research is isolating stem cells from other sources. It is one of the early but very important processes inherent in stem cell research. Subsequently, many methods have been developed to derive various kinds of stem cells, which depend on the cell they belong to and the tissue they originate from.

Embryonic Stem Cells (ESCs): The stem cells that are commonly used are the Embryonic Stem Cells, which are collected from the inner mass of the blastocysts, which are termed the early stages of the embryos after fertilization. This isolation method could freeze some in vitro procedures, such as immunosurgery or laser disaggregation, to purify the ICM from the trophectoderm (Xu et al., 2014). This is a very sensitive process because the ESCs which is supposed to be isolated

must be healthy and, indeed, pluripotent. Subsequently, they can be grown on conditioned MEFs or serum-free using knockout (KO) basal media. These media formulations usually include KSR, bFGF, and LIF, maintaining pluripotency and minimizing differentiation of ESCs (Lee, 2013).

Adult Stem Cells (ASCs): ASCs are in various tissues and organs of the adult organism, for example, the bone marrow, adipose tissue, and umbilical cord blood. The isolation methods of ASCs are not the same, depending on the tissue origin as well as the stem cell type. For example, blood-making stem cells known as hematopoietic stem cells (HSCs) are usually isolated from the bone marrow or the peripheral blood by methods. Such as density-gradient centrifugation or fluorescence-activated cell sorting, where cells are sorted depending on the receptors on their surface. MSCs capable of in vitro self-renewal and differentiating into multiple lineages while also exerting immunosuppressive effects (Bassi et al., 2011). These are obtained by either adherence of the culture to plastic or by immuno-selecting the culture for MSC markers from either the bone marrow aspirate.

Induced Pluripotent Stem Cells (iPSCs): The generation of iPSCs also involves the reprogramming of adult somatic cells like the skin fibroblasts, blood cells, etc., back to the state of embryonic-like cells using factors. This reprogramming process is made possible by the transduced four transcription factors: The genes Oct-4, Sox-2, Klf-4, and c-Myc to be delivered into the somatic cells (Jatving & Schepers). For this reason, practices like viral transduction, non-integrating vectors, and direct gene delivery, including proteins, or even mRNA, have been used. Subsequently, the iPSCs are maintained in the same way as ESCs to retain their ability to remain stem cells and not differentiate.

4.2 Stem Cells in the Culture

Due to tissue and organ generation, it is vital to organize stem cell long-term preservation and stem cells' cultivation in a culture medium for experimental and clinical needs. This has to be offered in sterile ways and conditions like culture media, adequate growth amenities, and types of substrates. Common culture media for both ESCs and iPSCs are based media combined with FBS or recognized substitutes for FBS since they supply the needed nutrients and growth factors essential for the cells' sustainable growth. Besides, to manifest the ability of cells to differentiate into multiple lineages, bFGF and LIF are added to the culture medium since they help retain the undifferentiated state of such cells (Shoni et al., 2014).




Figure 4. Stem Cell Isolation and Culture Method

Culture conditions of ASCs, including MSCs, are culture media containing FBS or hPL and specific growth factors and cytokine to support growth and maintain the ability of ASCs to differentiate into more than one cell lineage (Mangum et al., 2017). Therefore, specific cultural conditions may be necessary for various types of ASCs based on their characteristics and the conditions they specify. This is a feature that directly influences stem cells, this is in regards to the physical form of the culture system. Factors like oxygen tension, substrate stiffness, and cell density ways stem cell self-renewal, differentiation, and survival. For instance, low concentration of oxygen and high oxygen tension (hypoxia) enhances the self-renewal and survival of HSCs, and substrate stiffness controls the differentiation of MSCs into osteoblasts and adipocytes (X. Zhang

et al., 2022). Hence, it can be concluded that the stem cell isolation and culture methodology, or rather, their manipulation, has helped in the advancement of stem cell technology (Figure 4). The researchers are optimizing these techniques in every possible manner and the discovery of stem cell biology and realizing the potential in forming regenerating medicine and several other therapeutic methods.

4.3 Genetic and Epigenetic Modification

In addition, there is germ-line engineering and/or epigenetic modification that has advanced the control of stem cells and enhanced any genetically defective genes. These techniques have greatly improved the likelihood and strength of studying stem cells, disease simulation, and possible treatments.

CRISPR/Cas9 and other Gene Editing Tools

Clustered Regularly Interspaced Short Palindromic Repeats/CRISPR-associated protein 9 (CRISPR/Cas9) is an efficient and multimodal technique for stem cells' gene editing (Kato-Inui et al., 2018). A technique that allows for intentional genome modification by using double-strand breaks in the DNA that are then repaired either through NHEJ or HDR (Figure 5). NHEJ is relied on for gene disruption mostly in small insertions, while HDR is used in the generation either of single nucleotide variations (SNVs) or in s on inserting other sequences of interest in the genome. However, for the use of CRISPR/Cas9, has been used to genetically alter disorders in patient-specific iPSCs in stem cell research (Qin & Gao, 2016). For instance, some of the works revealed that the mutations associated with diseases such as cystic fibrosis, sickle cell anemia, and Duchenne muscular dystrophy can be repaired in iPSCs for autologous cell therapy. CRISPR/Cas9 has also been applied to introduce reporter genes and fluorescent tags into stem cells, tracking the behaviors and development as the stem cell differentiates (Bao et al., 2019). Hence, apart from the CRISPR/Cas9 system of gene editing, there are other modes of gene editing applied in stem cells, and they include the zinc finger nucleases (ZFNs) and the transcription activator-like effector nucleases (TALENs). These technologies utilize designed DNA binding proteins that affix to specific target sequences within the DNA to induce double-strand breaks that make gene editing quite easier, like that of CRISPR/Cas9 (Saber Sichani et al., 2023). Some of these technologies have been developed to some extent and implemented in practice, although less frequently than in the case of using recombinant proteins from the company's 'storehouse', because of the design and synthesis of the specific DNA-binding proteins required to control target genes.




Figure 5. Mechanism of Non-Homologous End-Joining of Double-Strand Breaks

Epigenetic modifications and Their effects

Epigenetic changes are also executed through DNA and histones as a means of DNA methylation to regulate the extent of gene expression rather than the actual genetic code. These changes are particularly reversible and can be controlled by various signals from the environment or developmental needs; that is why they are the subject of the study of stem cells (Shenghui et al., 2009). DNA methylation is commonly defined as an addition of a methyl group to the cytosine base of the DNA molecule and is typically observed in CpG dimer (Ren et al., 2018). This alteration is mostly associated with gene silencing and has a crucial role in maintaining the specialized characteristics of cells. For instance, stem cells exhibit demonstrably finite DNA methylation, and these can be rewired during the embryo organism's development and cell specialization process. Concerning the histones, the covalent modifications involve acetylation, methylation, phosphorylation, and ubiquitination, and these occur on histone proteins. It also carries the ability to alter the conformation or chromatin to give more or less access to the transcription factors and other regulating proteins to transcribe genes (Todeschini et al., 2014). Histone modifications are crucial for maintaining stem cells in an undifferentiated or pluripotent state and for altering between these two states, that is, the self-renewing state or the differentiation state of stem cells. Thus, considering the specifics of the epigenetic process of manipulating stem cells, it is possible to conclude that another tool for controlling cell fate and function has been revealed. For instance, the small molecules and other epigenomic factors for the differentiation of iPSCs to several cell types required for regenerative medicine are under investigation. Also, it is mentioned that epigenetic changes are linked with many diseases including cancer. Research concerning epigenetic events in stem cells can thus offer fresh management solutions to numerous diseases (Dompe et al., 2020).

4. 4. Stem Cell Imaging and Tracking

Consequently, cell labeling methodologies and imaging techniques have become firmly established as essential tools in stem cell biology to help researchers follow stem cell development within culture plates and animal models (Eglen & Reisine, 2011). They contribute to acquiring information about alterations in the behavior of stem cells, their migration, development, interaction with the adjacent cells, and their use in clinical procedures, which aids in setting up stem cell therapies in clinics.

Approaches to the monitoring of mesenchymal stem cells

Many methods have been employed in staining stem cells; several factors influence the efficiency of staining the particular stem cells depending on the intended application and the type of stem cells as well as the localization and time limitation of the chosen technique.

1. Genetic Labeling: This approach has used stem cells which have been engineered to express reporter genes, which include fluorescent proteins such as green fluorescent protein (GFP) or bioluminescent enzymes such as luciferase; hence, they can be seen over time with their progeny (Jurgielewicz et al., 2017). This method is flexible and can be achieved by viral transduction or transfection or the use of transgenic animal explants expressing the reporter gene placed under the regulation of the stem cell, particularly the promoter. Genetic labeling is used in long-term stem cell tracking it provides data concerning stem cell differentiation potential and tissue integration (Stroh et al., 2011). However, it involves changing the cells' genes, which can prove risky as far as clinical application is concerned.

2. Cell Surface Markers: Stem cells have certain markers, which are proteins attached to the cell surface and help recognize the cell type. These markers can be targeted by antibiotics tagged by fluorescent dyes with the help of labels; these markers can be visualized with flow cytometry or immunofluorescence microscopy (B. Zhang et al., 2022). This method is rather noninvasive and barely affects the cell, though alterations in cell surface marker expression density prevent long-term identification of certain cells.

3. Magnetic Resonance Imaging (MRI): MRI has in no way been known to be injurious to the human body, and it can be utilized to track stem cells that have been labeled with magnetic nanoparticles (Cromer Berman et al., 2011). Body cells then consume these particles and make a signal that can be checked through MRI machines. This allows one to track the stem cells in live animals and, therefore, get substantial data on biodistribution, migration, and engraftment in the tissues. However, to some of the other imaging techniques MRI could be comparatively less sensitive and lower in spatial resolution.

4. Nuclear Imaging: Albeit the radioactivity of the radioisotope used to label the stem cells could be technetium 99m or indium III, gamma rays are produced that can be identified using SPECT and or PET scanners (Bal, 2007). This procedure makes positron emission tomography possible and can be used to study the distribution of stem cells and their migration in the entire body without biopsy, but the technique requires the usage of radioactive materials, which is a complication concerning safety. Throughout the course of this dissertation, two broad approaches that aim to provide a more in-depth insight into the tissue, cell, and subcellular level characteristics of diseases will be described:

5. In vitro imaging: Stem cell imaging can be done on culture dishes or any man-made structures, such as under the microscope. Histo-to-chemical analysis and live cell imaging procedures are immensely useful for researchers to analyze stem cells' morphological, growth, and developmental status and their mobilization in the same context. These techniques are important to investigate the consequences of different conditions: other interventions such as drugs, cytokines, or genetic manipulation on the stem cell's activity (Ceccarelli et al., 2020).

6. In vivo imaging: This includes attempting to visualize the stem cells in their most natural habitats, which are the living organisms. Positive results attained by this knowledge include bioluminescence imaging, MRI, SPECT, and PET, which assist the researcher in observing stem cell transplantation in life to determine the survival and engraftment potential of cells and their differentiation ability in host tissue, in addition to therapeutic efficiency. Therefore, imaging is of significant use in stem cell-based therapies' evaluation both preclinically and in clinical settings in vivo (Klontzas et al., 2021).

5. Applications in Regenerative Medicine

It becomes one of the most complex tissues in the body mainly due to the neurons and glial cells that make it up, though they are believed to have low inherent regenerative capabilities. However, stem cell research has opened new chances for neural recovery for SCIs and NDDs, and these are very good developments in neural treatment.

5.1 Repairing Spinal Cord Injuries

Medical consequences of spinal cord lesions are paralysis and loss of feeling in the regions of the body beneath the area of the injury because the connection between the brain and the rest of the body is interrupted. The current stem cell therapies are said to unlock ways of enhancing processes in neural repair that affect SCI patients' overall functionality (Zhang et al., 2019).

Neural Stem Cells (NSCs): Labeled NSCs, as they are long-lived cells found in the adult mammalian, which are capable of self-renewal and differentiation into any cell of the nervous system including neurons, astrocytes, and oligodendrocytes; they are the three major classes of cells of the nervous system (Daynac & Petritsch, 2017). The preceding studies have shown that the propriety of NSCs for transplantation in the ischemic spinal cord can promote axonal regeneration, oligodendrocyte reforming, and functional recovery in animals. NSCs release factors that have therapeutic attributes on damaged neurons in the following manners: generate new healthy neurons and launch opinions that support the survival and growth of neurons besides decreasing inflammation. The current literature's research studies show that several clinical trials regarding

NSC transplantation are still undergoing for treating SCI patients (Hu et al., 2021).

Mesenchymal Stem Cells (MSCs): However, because MSCs are not neural stem cells, they have been reported to hold some potential in SCI, the outcome of which appears from immunomodulatory and neuroprotective properties. Studies have shown that MSC transplantation restores the blood barrier, decreases inflammation, enhances endothelial cell growth, remodeling, and repair in the injured cord, and, therefore, the functional outcomes in animal models (Lv et al., 2021). Clinical trials are also useful in establishing the benefits of MSCs in addressing situations of SCI patients.

5.2 Repairing Neurodegenerative Diseases

Neurodegenerative diseases, such as Parkinson's, Alzheimer's, or ALS, are diseases in which neurons and any functions have started fading. Critics have contended that stem cell therapies can replace the neurons that are destroyed, safeguard other neurons from the disease's effects, and slow down or halt the disease's progression.

Neural Stem Cells (NSCs): Neurogenic stem cell transplants have also been studied in the context of various NDDs in animal models, and the reports have revealed positive outcomes (Guerreiro & Maciel, 2023). Thus, Demetri and colleagues show that with the help of NSCs and the creation of dopaminergic neurons in the brains of animal models of Parkinson's disease, it is possible to enhance motor function. Research in animal models has since shown that NSC transplantation leads to a decrease in amyloid plaques and an increase in neurogenesis apart from the enhancement of cognition for Alzheimer's disease (Fouad, 2019).

Induced Pluripotent Stem Cells (iPSCs): The generation of iPSCs with the characteristics of the patient has opened up new vectors for apprehending and designing drugs for NDDs (Vijay et al., 2023). This is important because if you have iPSC-derived neurons carrying the genes for a specific disease, you can ascertain or determine the cause of the disease besides experimenting with drugs on specific neurons in a plate, a technique that is referred to as pharmacogenomics or, more specifically, personalized medicine. Besides, iPSC is meant for medical application with cell replacement therapies; this concerns the fact that neurons generated from patients' iPSCs may be grafted into the brain to replace the damaged ones (Paolini Sguazzi et al., 2021).

5.3 Neural Stem Cells: A New Development in the Field

For the ONS, discoveries that have come up have been of immense importance in studying the expansion and application of the theory. Certain of the recent tactics that have been developed were to refine the process of isolation and multiplication of NSCs and viability in transplantation (Abati et al., 2019). Hence, there are ongoing efforts to try and use biomaterials and tissue engineering to form structures through which stem cells that are to be transplanted, will be given direction and support as far as their functionality is concerned. Furthermore, technological advancements, particularly in gene editing like CRISPR/Cas9, have eased the chances of controlling NSCs for enhanced treatment (Lee et al., 2020). For instance, CRISPR/Cas9 has been used to cure NDDs based on fixing a mutational mistake in a couple of iPSC-derived neurons, The World's First Gene Editing for NDD treatment (De Guidi, 2021). However, some limitations have been raised in the current studies, which constitute a preliminary barrier to translating such results into practice. Further studies have to be conducted to determine the standards concerning stem cell transplants, raise the rates of survival and integration of the stem cells, and ascertain the efficiency and security of stem cell therapy.

5.4 Applications in Regenerative Medicine: Musculoskeletal Regeneration

Joint and muscular disorders Such disorders involve the bones, cartilage, or muscles and are prevalent hence being painful to the patient. Stem cells as a source of regenerative medicine can be

viewed as a perspective for repairing and regenerating these tissues; thus, the further development of orthopedic therapy with better chances to improve patient's quality of life may be provided (Akpancar et al., 2016).

Bone Repair

Most of the time, the bone heals itself; therefore, this process involves calling for help only when the defect is of a certain size or the healing process is in danger for one reason or another. Thus, the inherent characteristic of stem cells, particularly MSCs, to transform into osteoblasts makes stem cells provide substantial solutions to bone regeneration (Hollý et al., 2021). MSCs can be harvested from BM, Adipose tissue, and UC, and the cells can be either directly transplanted into the bone defect or placed into bioengineered scaffolds. The research done on pre-clinic and clinical studies has shown that MSC enhances bone repair in diseases such as nonunion fractures, osteonecrosis, and spinal fusion.

Cartilage Repair

Chondrocytes, which are the cells with the ability to produce cartilage, are relatively poor at repair and regeneration mechanisms even though cartilage is a smooth, flexible connective tissue that lays the slippery floor on all bones that rub together in the joints (Nochehdehi et al., 2023). As a result, cartilage pathologies are associated with such clinical manifestations as pain and stiffness and osteoarthritis as a form of degenerative-dystrophic changes in joints. The options for cartilage repair using stem cells include MSCs and chondrocytes or cartilage cells. MSCs can differentiate into chondrocytes and synthesize cartilage-specific ECM, therefore active in cartilage regeneration (Phull et al., 2016). Autologous chondrocyte implantation (ACI) involves the following procedures; first, the healthy cartilage cells are harvested from the patient. Next, those cells are cultured to make them look like healthy cells, and those cells are then implanted in the affected region. Some of the studies that have been conducted are as follows. ACI can be documented for reporting satisfactory results in the management of focal chondral and cartilaginous lesions in the knee (Jones & Peterson, 2007).

Muscle Repair

Such injuries tend to affect the basic structures and abilities of the affected body parts, with quality of life being seen more where muscles are involved. Among the cell-based therapies, some of the specific cells are muscle satellite cells and mesoangioblasts, which have the capability of muscle regeneration. These stem cells are present either between the muscle fibers or as a component of the nuclei of the muscle cells; their main functions are muscle hypertrophy and regeneration (Blaauw & Reggiani, 2014). The Mesoangioblasts are the kind of stem cells that are vessel-associated and dwelling in skeletal muscles; they can pen various kinds of cells with the help of fusion through which muscular fibers are also produced (Yedigaryan et al., 2022). Ex vivo, the future of these stem cells has demonstrated their ability to enhance muscle repair upon injury and boost strength and functioning (Qazi et al., 2019).

5.5 Applications in Orthopedic Treatments

Stem cell-based therapies are being investigated for a wide range of orthopedic applications, including research in stem cell applications for orthopedics has been active for numerous applications such as:

Osteoarthritis: MSCs and other stem cell types are discussed for the limitation or radical cure of OA by promoting cartilage regeneration and modulating the inflammation and immune responses (Liu et al., 2022).

Non-union fractures: MSCs, along with BMAC, consist of a mixture of stem and progenitor cells and are used to enhance bone formation in non-union fractures due to the body's inability to

heal (Haeusner et al., 2023).

Tendon and ligament injuries: At the time, many different stem cells, such as TDSCs and MSCs, have been investigated, and the tendon and ligament healing potentiality (Wei & Lu, 2021) of these cells has been studied for improving post-injurious healing and reducing the probability of re-injury. However, certain problems, such as the optimal way of delivering stem cells, the source of a constant stem cell supply for the patients, and the other effects of these anti-osteoporotic therapies in the perioperative period of the surgery, have not been optimized yet. Such modern approaches are focused on solving these problems and transferring the possible outcomes of preclinical investigations into clinical applications for managing musculoskeletal diseases.

5. 7 Applications in Regenerative Medicine: hematopoietic stem cell transplantation (HSCT)

Nevertheless, hematopoietic stem cell transplantation (HSCT) is the possible curative treatment method concerning the capabilities of hematopoietic stem cells (HSCs) to provide the repair of impaired or mutilated bone marrow (Alvin, 2021). HSCs are found in the BM, PB, and cord blood they are, by default, the only cells that make all the blood cells and are, therefore, ideal for re-populating BM of patients with diseased blood cell-making factories.

Treatments arise from the assessment of blood-related diseases and cancer genes

HSCT has, however, been useful in the management of a combination of hematological malignancies, for example, leukemia, lymphoma, and multiple myelomas (A Patel & Rameshwar, 2011). HSCT is also applied in managing other nonneoplastic hematological diseases like aplastic anemia, sickle cell anemia, and thalassemia. HSCT can be either autologous, in which harvested Stem cells are used from the patient's/own marrow and Stem cells, which are harvested from the Stem cells of a related donor, or a matched unrelated donor and syngeneic Stem cells harvested from an identical twin. In the context of cancer treatment, HSCT serves two main purposes in which concerning cancer therapy, HSCT has two principal functions:

1. Myeloablative Therapy: To treat Hepatoid carcinoma, the patient is given a high intensity of chemotherapy or radiation to eliminate the cancerous cells in the bone marrow. While this treatment eliminates the body's required, healthy HSCs tend to myelosuppression as well as make the patient immune deficient (DeFilipp et al., 2022).

2. Stem Cell Rescue: HSCs are transplanted by replacing the defective hematopoietic system of the body either from the patient's bone marrow or from donor bone marrow, which is referred to as allogeneic transplant in cases where the patient's transplant, which is referred to as autologous transplant is not feasible. The HSCs are introduced in the bone marrow of the patients, where they differentiate into different kinds of blood cells, providing a direct replacement to the unhealthy cells or those that may be invaded by the disease (Kandarakov et al., 2022). In the case of nonmalignant hematologic diseases, the procedure for HSCT involves transplanting normal HSCs into the patients' bodies to replenish normal blood and modify the character of hematological disorders.

Current State and Challenges in Organ Transplantation

HSCT has evolved over the years owing to the techniques of stem cell mobilization, collection, and processing, and changes in the condition for pre-transplant regimens, which has improved survival and less toxicity (Wade, 2023). Another difference is the increase in the use of peripheral blood stem cells (PBSCs) compared to bone marrow because it is easy to harvest and engraft faster. There have also been non-myeloablative or reduced-intensity preparations made for elderly patients and patients with previous illnesses; therefore, HSCT can be applied to more patients (Lekakis & De Lima, 2008). However, this field still poses some difficulties in the case of an increase in the complexity of HSCT and potential complications, including GVHD, infection, and organ toxicity

(Reis et al., 2016). The problem arises where the conditioning regimen of the donor stem cells recognizes the recipient's tissues as foreign entities and proceeds to produce immune cells to attack the target tissue, which may range from mild form to severe form. Morbidity concerns result from immunosuppression, which is witnessed after the conditioning regimen; acute organ toxicity tendencies result from either the high dose chemotherapy or radiation therapy (Weichselbaum et al., 1980). Recovery research today entails efforts to minimize the mentioned complications; this involves donor-recipient compatibility, choice of regimen conditioning, and measures of combating GVHD (Timofeeva et al., 2022). New ways of promoting the culture of HSCs and engraftment are also being established to enhance the efficiency of the HSCT. Hematopoietic stem cell transplantation remains one of the key advancements in managing blood disorders and malignancy, as it offers definitive therapy to many of the affected clients. That is why new advancements in transplantation surgery, along with advancements in the management of such patients, have effectively improved the salvage modality of this intervention.

6. Roles in Disease Simulation and New Drugs Development

Due to the development of stem cells, especially iPSCs, disease modeling and drug discovery have been enhanced. Techniques of deriving the patient's iPSCs and their subsequent differentiation into multiple lineages have provided new roads toward understanding the pathogenesis of diseases, identifying biomarkers, and finding specific therapies for patients.

Disease Modeling

One serious deficiency of the present disease models is that they are either animal models or immortalized cell lines and do not represent human diseases regarding genetics and phenotype. Unlike this, iPSCs can be derived exclusively from patients with specific diseases with clearly identifiable genetic mutations and epigenetic tags identical to the patient's cells (Jang & Ye, 2016). This makes it possible to create disease models that replicate the human organism's status as it assists in understanding diseases' mechanisms and pathogenesis. For instance, iPSCs derived from patients with neurodegenerative diseases, Parkinson's and Alzheimer's disease, have been utilized in modeling such diseases and understanding the molecular activity and cellular signaling routes (Yefroyev & Jin, 2022). In this manner, it is possible to investigate the neuronal differentiation of iPSC, the appearance of a disease, or certain processes, such as the formation of aggregates and neuronal death in a plate. These models have assisted in identifying new therapies for all these diseases besides testing compounds that could assist in preventing or treating these diseases. Similarly, iPSCs of patients with Cardiomyopathy and Arrhythmia have been applied to fashion in vitro diseases of the Cardiovascular system and to compare the cellular phenotypes consequent to disease-relevant alleles. These models have assisted in elucidating the processes of cardiac dysfunctions and in the development of new drugs that operate at molecular levels (McGregor & Dunn, 2006).

Drug Screening and Toxicology

Thus, the stem cell-based disease models are beneficial in drug discovery and also in assessing the toxic effects. An existing technique in drug identification largely relies on animal testing, and success attained from such could be far from an actual estimate of successful drugs in man because of species differences (Liebsch et al., 2011). On the other hand, the cells derived from iPSCs represent a real human/system of speaking to evaluate the response to drugs and possible adverse effects. Thus, the iPSC-based HTS platforms for drug discovery have been developed so that a large number of compounds in the library can be tested for therapeutic potential. It can also identify small molecules that may preserve disease phenotypes, repair diseased cells, or modulate specific signaling pathways. There are also opportunities to compare the differences in drug sensitivity in the iPSC-derived cells originating from patients with different genetic profiles or disease types (Grskovic et al., 2011). Furthermore, the derived cells from the iPSCs can be used for the study of

the toxicity of the desired drug candidate on the stem cell's viability, proliferation, differentiation, and functional profile alteration (Genova et al., 2019). This makes it possible for the researchers to identify some of the safety problems before the drug is ready to be injected into patients, which goes a long way in preventing the impacts on the patients participating in clinical trials.

7. Future Directions and Recommendations

Stem cell research is a relatively new and rapidly developing branch of science in which discoveries can dramatically alter the functioning of medicine. However, some issues and ethical factors are also taken into consideration regarding the intended objectives for applying stem cell-based therapy.

Enhancing stem cell survival and engraftment: One of the major challenges of stem cell transplants is that cells do not fare very well because engraftment is fairly poor. The current strategy is to find a definitive mode of handling the problem of survival and integration of the transplanted cell into the host tissue; this uses ideas concerning biomaterials as well as tissue engineering (Mooranian et al., 2021).

Optimizing stem cell differentiation and maturation: Nevertheless, it is difficult to continue to possess the abilities of differentiation to various cell types, direct their differentiation, and substantiate that they become functional cells. Research should be done concerning the type of signals that promote the differentiation of stem cells into the desirable lineages and the culture conditions that promote the differentiation of stem cells into fully mature cells (Cai et al., 2007).

Addressing safety concerns: Even in cell therapy, the stem cells that are prospective for the treatment of diseases are still able to form tumors or are recognized as foreign bodies by the immune system of the body. The long-term safety of stem cell therapies also poses challenges for preclinical and clinical trials (Squillaro et al., 2016). It must be performed before the recipients' immune systems can be protected against harm resulting from the erroneous placement of these cells and their migration to locations that are problematic for the recipient's body.

Scaling up production and ensuring quality control: Because stem cell clinical application requires mass production of stem cells, this can only be achieved in a process that affords cell quality and consistency in the stem cell manufacturing process. Therefore, enhancing quality assurance and developing policies concerning stem cells is important to apply such findings to practice (Papassavas et al., 2015).

Personalizing stem cell therapies: Popular information technologies such as genomic and Precision medicine have provided options for personal stem cell therapies. They include disease/Genetic Map responsive treatment. This strategy can also raise the efficiency of stem cell therapies because it identifies the prospect of a specific patient reacting positively to a treatment and minimizes the chances for an adverse outcome (Desgres & Menasche, 2019). Recommendations for future research and development include:

1. Investing in basic research: Stem cell therapies as safe, effective means of treatment, additional insight into stem cell biology, namely the mechanisms of stem cell self-renewal and determination as well as their interactions with the immediate environment, is needed (Li et al., 2022).

2. Developing standardized protocols: Another requirement is the severity of procedures applied to stem cells' isolation as well as their further cultivation and differentiation within cells, as this condition influences the possibility of achieving the same results in different labs or clinics (Sensebe et al., 2010).

3. Collaborating across disciplines: Stem cell research is an area of research that means that various professionals, engineers, biologists, clinicians, and regulatory scientists can implement

what has been learned in the laboratories (Wainwright et al., 2006).

4. Investing in education and outreach: Stem cell therapy is also an unclear area, so society and those offering care need to educate themselves on the positive changes that may result from the use of new technology and the dangers attributed to stem cell therapies (Einsiedel & Adamson, 2012). Due to stem cells' ability of future changes in the medical practice and millions of patients, it can be considered as one of the most significant discoveries. If one comprehends and manages to combat the difficulties and ethical issues arising in the course of completing the work of this chapter, advancing the efforts in researching, developing, and enlightening, one can contribute to the potential of stem cell therapies and open the possibility to enhance the future of the regenerative medicine.

8. Conclusion

Stem cell science has become one of the most promising fields in modern health care science, offering new directions in understanding diseases, treating them, and even regenerating tissues. Because of their properties, as well as their ability to replenish and convert themselves into any cell of choice, stem cells apply to creating new therapeutic models. In this chapter, researchers have differentiated stem cells in brief by classification, such as embryonic stem cells, adult stem cells, and induced pluripotent stem cells, and have tried to exhibit the picture regarding the differences and the features of each kind of stem cell broadly considering the possibilities of their use in the treatment of diseases. Two of them explained how stem cells operate and the factors, including stem cell microenvironments, signaling, and epigenetic mechanisms controlling self-renewal, differentiation, and plasticity. Furthermore, we have discussed the future application of stem cells in regeneration medicine, such as cardiovascular regeneration, neural regeneration, musculoskeletal tissue regeneration, and Hematopoietic stem cell transplantation. We have also mentioned using stem cells in disease modeling, which means disease and drug analysis and screening, as well as customized medicine. In total, the successes of stem cell research are quite sensational; nonetheless, the challenges remain critical. Stem cell-based therapies' safety and effectiveness, how the cells are delivered how the cells engraft and the ethics of the issues all are crucial in the attempt to translate discoveries in the lab to real-life scenarios. For this; as stem cell biology, engineering and the successful translation of such findings into medical applications progresses, stem cell-based therapies should have a promising future. Thus, stem cell research can be concluded to be a significant discovery in medical science that has transformed the very picture of the modern curing paradigm and, at the same time, opened up brand new vistas for the treatment of countless diseases. When proper respect is accorded to stem cells, human beings can advance the usefulness for reshaping tilted tissues and severe diseases and, in this way, boost health. The road might be full of challenges; however, the reward is priceless; thus, come up with the vision of a future where regenerative medicine will be the backbone of medicine.

References

- A Patel, S., & Rameshwar, P. (2011). Stem cell transplantation for hematological malignancies: prospects for personalized medicine and co-therapy with mesenchymal stem cells. *Current Pharmacogenomics and Personalized Medicine (Formerly Current Pharmacogenomics)*, 9(3), 229-239.
- Abati, E., Bresolin, N., Comi, G. P., & Corti, S. (2019). Preconditioning and cellular engineering to increase the survival of transplanted neural stem cells for motor neuron disease therapy. *Molecular Neurobiology*, 56, 3356-3367.
- Akpancar, S., Tatar, O., Turgut, H., Akyildiz, F., & Ekinici, S. (2016). The current perspectives of stem cell therapy in orthopedic surgery. *Archives of trauma research*, 5(4).
- Al-Jedai, A. H. (2008). Notch signalling in CD34+ cells in chronic myeloid leukaemia. *The*

University of Manchester (United Kingdom).

Alvin, G. (2021). Cell therapy: the new approach to der-matology and dermatologic surgery. *Clin Surg*, 5(8), 1-14.

Baba, A. I., & Cătoi, C. (2007). Tumor cell morphology. In *Comparative oncology*. The Publishing House of the Romanian Academy.

Bal, C. (2007). Radio-isotopic techniques. In.

Bao, X., Adil, M. M., Muckom, R., Zimmermann, J. A., Tran, A., Suhy, N., Xu, Y., Sampayo, R. G., Clark, D. S., & Schaffer, D. V. (2019). Gene editing to generate versatile human pluripotent stem cell reporter lines for analysis of differentiation and lineage tracing. *Stem Cells*, 37(12), 1556-1566.

Bassi, Ê. J., Aita, C. A. M., & Câmara, N. O. S. (2011). Immune regulatory properties of multipotent mesenchymal stromal cells: where do we stand? *World Journal of Stem Cells*, 3(1), 1.

Beerman, I., & Rossi, D. J. (2015). Epigenetic control of stem cell potential during homeostasis, aging, and disease. *Cell stem cell*, 16(6), 613-625.

Birbrair, A., & Frenette, P. S. (2016). Niche heterogeneity in the bone marrow. *Annals of the New York Academy of Sciences*, 1370(1), 82-96.

Blaauw, B., & Reggiani, C. (2014). The role of satellite cells in muscle hypertrophy. *Journal of muscle research and cell motility*, 35, 3-10.

Bonaventura, G., Iemmolo, R., Attaguile, G. A., La Cognata, V., Pistone, B. S., Raudino, G., D'Agata, V., Cantarella, G., Barcellona, M. L., & Cavallaro, S. (2021). iPSCs: A preclinical drug research tool for neurological disorders. *International journal of molecular sciences*, 22(9), 4596.

Burchfield, J. S., & Dimmeler, S. (2008). Role of paracrine factors in stem and progenitor cell mediated cardiac repair and tissue fibrosis. *Fibrogenesis & tissue repair*, 1, 1-11.

Burel, M. (2022). *Microenvironment Remodeling as a Novel Mechanism for Germline Stem Cell Competition* [New York University].

Cai, L., Ye, Z., Zhou, B. Y., Mali, P., Zhou, C., & Cheng, L. (2007). Promoting human embryonic stem cell renewal or differentiation by modulating Wnt signal and culture conditions. *Cell research*, 17(1), 62-72.

Ceccarelli, S., Pontecorvi, P., Anastasiadou, E., Napoli, C., & Marchese, C. (2020). Immunomodulatory effect of adipose-derived stem cells: the cutting edge of clinical application. *Frontiers in Cell and Developmental Biology*, 8, 236.

Çetin, E. A., Babayiğit, E. H., Özdemir, A. Y., Erfen, Ş., & Onur, M. A. (2023). Investigation of UV-treated mesenchymal stem cells in an in vitro wound model. *In Vitro Cellular & Developmental Biology-Animal*, 59(5), 331-345.

Cromer Berman, S. M., Walczak, P., & Bulte, J. W. (2011). Tracking stem cells using magnetic nanoparticles. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 3(4), 343-355.

Dawson, R., Beal, M. F., Bondy, S. C., DiMonte, D., & Isom, G. E. (1995). Excitotoxins, aging, and environmental neurotoxins: implications for understanding human neurodegenerative diseases. *Toxicology and applied pharmacology*, 134(1), 1-17.

Daynac, M., & Petritsch, C. K. (2017). Regulation of asymmetric cell division in mammalian neural stem and cancer precursor cells. *Asymmetric Cell Division in Development, Differentiation and Cancer*, 375-399.

- De Guidi, C. (2021). Generation of induced pluripotent stem cells (iPSCs) lines deficient for genes associated with neurodevelopmental diseases using CRISPR/Cas9 technology. In.
- DeFilipp, Z., Hefazi, M., Chen, Y.-B., & Blazar, B. R. (2022). Emerging approaches to improve allogeneic hematopoietic cell transplantation outcomes for nonmalignant diseases. *Blood, The Journal of the American Society of Hematology*, 139(25), 3583-3593.
- Desgres, M., & Menasche, P. (2019). Clinical translation of pluripotent stem cell therapies: challenges and considerations. *Cell stem cell*, 25(5), 594-606.
- Desmarais, J. A., Demers, S.-P., Suzuki, J., Laflamme, S., Vincent, P., Laverty, S., & Smith, L. C. (2011). Trophoblast stem cell marker gene expression in inner cell mass-derived cells from parthenogenetic equine embryos. *Reproduction*, 141(3), 321.
- Dompe, C., Janowicz, K., Hutchings, G., Moncrieff, L., Jankowski, M., Nawrocki, M. J., Józłowski, M., Mozdziak, P., Petite, J., & Shibli, J. A. (2020). Epigenetic research in stem cell bioengineering—anti-cancer therapy, regenerative and reconstructive medicine in human clinical trials. *Cancers*, 12(4), 1016.
- Eglen, R., & Reisine, T. (2011). Primary cells and stem cells in drug discovery: emerging tools for high-throughput screening. *Assay and drug development technologies*, 9(2), 108-124.
- Einsiedel, E. F., & Adamson, H. (2012). Stem cell tourism and future stem cell tourists: policy and ethical implications. *Developing world bioethics*, 12(1), 35-44.
- Ejtehadifar, M., Shamsasenjan, K., Movassaghpour, A., Akbarzadehlaleh, P., Dehdilani, N., Abbasi, P., Molaeipour, Z., & Saleh, M. (2015). The effect of hypoxia on mesenchymal stem cell biology. *Advanced pharmaceutical bulletin*, 5(2), 141.
- Ericson, R. J. (2007). Bridging solutions to the religion and science conflict over human embryonic stem cell* research. George Mason University.
- Foster, C. S., Dodson, A., Karavana, V., Smith, P., & Ke, Y. (2002). Prostatic stem cells. *The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland*, 197(4), 551-565.
- Fouad, G. I. (2019). Stem cells as a promising therapeutic approach for Alzheimer's disease: a review. *Bulletin of the National Research Centre*, 43(1), 52.
- Genova, E., Cavion, F., Lucafò, M., De Leo, L., Pelin, M., Stocco, G., & Decorti, G. (2019). Induced pluripotent stem cells for therapy personalization in pediatric patients: focus on drug-induced adverse events. *World Journal of Stem Cells*, 11(12), 1020.
- Gordeeva, O. (2019). TGF β family signaling pathways in pluripotent and teratocarcinoma stem cells' fate decisions: balancing between self-renewal, differentiation, and cancer. *Cells*, 8(12), 1500.
- Grskovic, M., Javaherian, A., Strulovici, B., & Daley, G. Q. (2011). Induced pluripotent stem cells—opportunities for disease modelling and drug discovery. *Nature Reviews Drug Discovery*, 10(12), 915-929.
- Guerreiro, S., & Maciel, P. (2023). Transition from animal-based to human induced pluripotent stem cells (iPSCs)-based models of neurodevelopmental disorders: Opportunities and challenges. *Cells*, 12(4), 538.
- Haeusner, S., Jauković, A., Kupeczyk, E., & Herrmann, M. (2023). cellularity in bone marrow autografts for bone and fracture healing. *American Journal of Physiology-Cell Physiology*, 324(2), C517-C531.

- Hall, P. A., & Watt, F. M. (1989). Stem cells: the generation and maintenance of cellular diversity. *Development*, 106(4), 619-633.
- Hollý, D., Klein, M., Mazreku, M., Zamborský, R., Polák, Š., Danišovič, L., & Csöbönyeiová, M. (2021). Stem Cells and Their Derivatives—Implications for Alveolar Bone Regeneration: A Comprehensive Review. *International journal of molecular sciences*, 22(21), 11746.
- Hu, X.-C., Lu, Y.-B., Yang, Y.-N., Kang, X.-W., Wang, Y.-G., Ma, B., & Xing, S. (2021). Progress in clinical trials of cell transplantation for the treatment of spinal cord injury: how many questions remain unanswered? *Neural regeneration research*, 16(3), 405-413.
- Ikegami, K., Ohgane, J., Tanaka, S., Yagi, S., & Shiota, K. (2009). Interplay between DNA methylation, histone modification and chromatin remodeling in stem cells and during development. *International Journal of Developmental Biology*, 53.
- Jang, Y.-Y., & Ye, Z. (2016). Gene correction in patient-specific iPSCs for therapy development and disease modeling. *Human genetics*, 135(9), 1041-1058.
- Jatving, M., & Schepers, H. Induced pluripotent stem cells: Wilt they be safe?
- Jayaraman, S., Rajagopal, P., Periyasamy, V., Palaniyandi, K., Kumaran, R. I., Reddy, S. V., Balasubramanian, S., & Sambandam, Y. (2021). Signaling pathways influencing stem cell self-renewal and differentiation—special emphasis on cardiomyocytes. In *Stem cells and aging* (pp. 157-168). Elsevier.
- Jha, B. S., Farnoodian, M., & Bharti, K. (2021). Regulatory considerations for developing a phase I investigational new drug application for autologous induced pluripotent stem cells-based therapy product. *Stem Cells Translational Medicine*, 10(2), 198-208.
- Jones, D. G., & Peterson, L. (2007). Autologous chondrocyte implantation. *Cartilage repair strategies*, 137-165.
- Jurgielewicz, P., Harmsen, S., Wei, E., Bachmann, M. H., Ting, R., & Aras, O. (2017). New imaging probes to track cell fate: reporter genes in stem cell research. *Cellular and Molecular Life Sciences*, 74, 4455-4469.
- Kandarakov, O., Belyavsky, A., & Semenova, E. (2022). Bone marrow niches of hematopoietic stem and progenitor cells. *International journal of molecular sciences*, 23(8), 4462.
- Karpowicz, P. A. (2004). A scientific and ethical investigation into the human retinal stem cell chimera: Manufacturing a monster? National Library of Canada= Bibliothèque nationale du Canada, Ottawa.
- Kato-Inui, T., Takahashi, G., Hsu, S., & Miyaoka, Y. (2018). Clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein 9 with improved proof-reading enhances homology-directed repair. *Nucleic acids research*, 46(9), 4677-4688.
- Khan, F. A., Almohazey, D., Alomari, M., & Almoftly, S. A. (2018). Isolation, culture, and functional characterization of human embryonic stem cells: current trends and challenges. *Stem cells international*, 2018(1), 1429351.
- Klontzas, M. E., Kakkos, G. A., Papadakis, G. Z., Marias, K., & Karantanis, A. H. (2021). Advanced clinical imaging for the evaluation of stem cell based therapies. *Expert Opinion on Biological Therapy*, 21(9), 1253-1264.
- Lee, J., Bayarsaikhan, D., Bayarsaikhan, G., Kim, J.-S., Schwarzbach, E., & Lee, B. (2020). Recent advances in genome editing of stem cells for drug discovery and therapeutic application. *Pharmacology & therapeutics*, 209, 107501.

- Lee, K.-H. (2013). Conditions and techniques for mouse embryonic stem cell derivation and culture. *Pluripotent stem cells*, 1, 85-116.
- Lekakis, L., & De Lima, M. (2008). Reduced-intensity conditioning and allogeneic hematopoietic stem cell transplantation for acute myeloid leukemia. *Expert review of anticancer therapy*, 8(5), 785-798.
- Lewandowski, J., & Kurpisz, M. (2016). Techniques of human embryonic stem cell and induced pluripotent stem cell derivation. *Archivum immunologiae et therapiae experimentalis*, 64, 349-370.
- Li, M., Jiang, Y., Hou, Q., Zhao, Y., Zhong, L., & Fu, X. (2022). Potential pre-activation strategies for improving therapeutic efficacy of mesenchymal stem cells: current status and future prospects. *Stem cell research & therapy*, 13(1), 146.
- Liebsch, M., Grune, B., Seiler, A., Butzke, D., Oelgeschläger, M., Pirow, R., Adler, S., Riebeling, C., & Luch, A. (2011). Alternatives to animal testing: current status and future perspectives. *Archives of toxicology*, 85, 841-858.
- Liu, S., Deng, Z., Chen, K., Jian, S., Zhou, F., Yang, Y., Fu, Z., Xie, H., Xiong, J., & Zhu, W. (2022). Cartilage tissue engineering: From proinflammatory and anti-inflammatory cytokines to osteoarthritis treatments. *Molecular Medicine Reports*, 25(3), 1-15.
- Lloyd-Lewis, B., Mourikis, P., & Fre, S. (2019). Notch signalling: sensor and instructor of the microenvironment to coordinate cell fate and organ morphogenesis. *Current opinion in cell biology*, 61, 16-23.
- Lv, B., Zhang, X., Yuan, J., Chen, Y., Ding, H., Cao, X., & Huang, A. (2021). Biomaterial-supported MSC transplantation enhances cell–cell communication for spinal cord injury. *Stem cell research & therapy*, 12, 1-16.
- Malgieri, A., Kantzari, E., Patrizi, M. P., & Gambardella, S. (2010). Bone marrow and umbilical cord blood human mesenchymal stem cells: state of the art. *International journal of clinical and experimental medicine*, 3(4), 248.
- Mangum, L. H., Natesan, S., Stone, R., Wrice, N. L., Larson, D. A., Florell, K. F., Christy, B. A., Herzig, M. C., Cap, A. P., & Christy, R. J. (2017). Tissue source and cell expansion condition influence phenotypic changes of adipose-derived stem cells. *Stem cells international*, 2017(1), 7108458.
- McGregor, E., & Dunn, M. J. (2006). Proteomics of the heart: unraveling disease. *Circulation research*, 98(3), 309-321.
- Mong, J. (2013). Integrating extrinsic and intrinsic cues to guide cell fate decisions: rational approaches in stem cell engineering. *Karolinska Institutet (Sweden)*.
- Mooranian, A., Jones, M., Ionescu, C. M., Walker, D., Wagle, S. R., Kovacevic, B., Chester, J., Foster, T., Johnston, E., & Kuthubutheen, J. (2021). Artificial cell encapsulation for biomaterials and tissue bio-nanoengineering: history, achievements, limitations, and future work for potential clinical applications and transplantation. *Journal of Functional Biomaterials*, 12(4), 68.
- Moses, B. S. (2016). The role of miRNAs in niche regulation of ALL phenotype. *West Virginia University*.
- Nochehdehi, A. R., Nemavhola, F., Thomas, S., & Maria, H. J. (2023). *Cartilage Tissue and Knee Joint Biomechanics: Fundamentals, Characterization and Modelling*. Elsevier.
- Paolini Sguazzi, G., Muto, V., Tartaglia, M., Bertini, E., & Compagnucci, C. (2021). Induced

pluripotent stem cells (iPSCs) and gene therapy: a new era for the treatment of neurological diseases. *International journal of molecular sciences*, 22(24), 13674.

Papassavas, A., Chatzistamatiou, T. K., Michalopoulos, E., Serafetinidi, M., Gkioka, V., Markogianni, E., & Stavropoulos-Giokas, C. (2015). Quality management systems including accreditation standards. In *Cord Blood Stem Cells and Regenerative Medicine* (pp. 229-248). Elsevier.

Phull, A.-R., Eo, S.-H., Abbas, Q., Ahmed, M., & Kim, S. J. (2016). Applications of chondrocyte-based cartilage engineering: an overview. *BioMed research international*, 2016(1), 1879837.

Portale, F. (2018). Activin A as a new key factor in the leukemic bone marrow niche.

Qazi, T. H., Duda, G. N., Ort, M. J., Perka, C., Geissler, S., & Winkler, T. (2019). Cell therapy to improve regeneration of skeletal muscle injuries. *Journal of cachexia, sarcopenia and muscle*, 10(3), 501-516.

Qin, Y., & Gao, W.-Q. (2016). Concise review: patient-derived stem cell research for monogenic disorders. *Stem Cells*, 34(1), 44-54.

Ranjitha, B. (2019). A Study of Haematological Parameters and Bone Marrow Morphology of Pancytopenia in Adult Patients Rajiv Gandhi University of Health Sciences (India)].

Reis, M., Ogonek, J., Qesari, M., Borges, N. M., Nicholson, L., Preußner, L., Dickinson, A. M., Wang, X.-n., Weissinger, E. M., & Richter, A. (2016). Recent developments in cellular immunotherapy for HSCT-associated complications. *Frontiers in immunology*, 7, 500.

Ren, R., Horton, J. R., Zhang, X., Blumenthal, R. M., & Cheng, X. (2018). Detecting and interpreting DNA methylation marks. *Current opinion in structural biology*, 53, 88-99.

Rhind, N., & Russell, P. (2012). Signaling pathways that regulate cell division. *Cold Spring Harbor perspectives in biology*, 4(10), a005942.

Ringe, J., Kaps, C., Burmester, G.-R., & Sitterling, M. (2002). Stem cells for regenerative medicine: advances in the engineering of tissues and organs. *Naturwissenschaften*, 89(8), 338-351.

Saber Sichani, A., Ranjbar, M., Baneshi, M., Torabi Zadeh, F., & Fallahi, J. (2023). A review on advanced CRISPR-based genome-editing tools: base editing and prime editing. *Molecular Biotechnology*, 65(6), 849-860.

Saltzman, W. M. (2004). *Tissue engineering: engineering principles for the design of replacement organs and tissues*. Oxford university press.

Šarić, T., Mehrjardi, N. Z., & Hescheler, J. (2010). Alternative Embryonic Stem Cell Sources. *Stem Cell Biology in Health and Disease*, 101-143.

Schewe, M., & Fodde, R. (2018). Multitasking Paneth cells in the intestinal stem cell niche. In *Advances in Stem Cells and their Niches* (Vol. 2, pp. 41-75). Elsevier.

Schwartz, S. D., Tan, G., Hosseini, H., & Nagiel, A. (2016). Subretinal transplantation of embryonic stem cell-derived retinal pigment epithelium for the treatment of macular degeneration: an assessment at 4 years. *Investigative ophthalmology & visual science*, 57(5), ORSFC1-ORSFC9.

Sensebe, L., Krampera, M., Schrezenmeier, H., Bourin, P., & Giordano, R. (2010). Mesenchymal stem cells for clinical application. *Vox Sanguinis*, 98(2), 93-107.

Shenghui, H., Nakada, D., & Morrison, S. J. (2009). Mechanisms of stem cell self-renewal. *Annual Review of Cell and Developmental*, 25(1), 377-406.

Shoni, M., O Lui, K., G Vavvas, D., G Muto, M., S Berkowitz, R., Vlahos, N., & Ng, S.-W.

(2014). Protein kinases and associated pathways in pluripotent state and lineage differentiation. *Current stem cell research & therapy*, 9(5), 366-387.

Sidrat, T., Rehman, Z.-U., Joo, M.-D., Lee, K.-L., & Kong, I.-K. (2021). Wnt/ β -catenin pathway-mediated PPAR δ expression during embryonic development differentiation and disease. *International journal of molecular sciences*, 22(4), 1854.

Singh, A., Yadav, C., Tabassum, N., Bajpeyee, A., & Verma, V. (2019). Stem cell niche: Dynamic neighbor of stem cells. *European journal of cell biology*, 98(2-4), 65-73.

Sokol, S. Y. (2011). Maintaining embryonic stem cell pluripotency with Wnt signaling. *Development*, 138(20), 4341-4350.

Sordi, V., Pellegrini, S., Krampera, M., Marchetti, P., Pessina, A., Ciardelli, G., Fadini, G., Pintus, C., Pantè, G., & Piemonti, L. (2017). Stem cells to restore insulin production and cure diabetes. *Nutrition, Metabolism and Cardiovascular Diseases*, 27(7), 583-600.

Sorensen, J. R., Mcfaline-Figueroa, J., & Call, J. A. (2022). Pathophysiology of volumetric muscle loss and targets for regenerative rehabilitation. In *Regenerative rehabilitation: From basic science to the clinic* (pp. 177-225). Springer.

Soria, B., Skoudy, A., & Martin, F. (2001). From stem cells to beta cells: new strategies in cell therapy of diabetes mellitus. *Diabetologia*, 44, 407-415.

Squillaro, T., Peluso, G., & Galderisi, U. (2016). Clinical trials with mesenchymal stem cells: an update. *Cell transplantation*, 25(5), 829-848.

Stroh, A., Tsai, H.-C., Wang, L.-P., Zhang, F., Kressel, J., Aravanis, A., Santhanam, N., Deisseroth, K., Konnerth, A., & Schneider, M. B. (2011). Tracking stem cell differentiation in the setting of automated optogenetic stimulation. *Stem Cells*, 29(1), 78-88.

Suter, D. M. (2020). Transcription factors and DNA play hide and seek. *Trends in cell biology*, 30(6), 491-500.

Timofeeva, O. A., Philogene, M. C., & Zhang, Q. J. (2022). Current donor selection strategies for allogeneic hematopoietic cell transplantation. *Human Immunology*, 83(10), 674-686.

Todeschini, A.-L., Georges, A., & Veitia, R. A. (2014). Transcription factors: specific DNA binding and specific gene regulation. *Trends in genetics*, 30(6), 211-219.

Uzbas, F., May, I., Parisi, A., Thompson, S., Kaya, A., Perkins, A., & Memili, E. (2015). Molecular physiognomies and applications of adipose-derived stem cells. *Stem cell reviews and reports*, 11, 298-308.

Verma, D. P. S., & Chatterjee, J. (2009). TORing with cell cycle, nutrients, stress and growth. In (pp. 161-180): Wiley Online Library.

Vijay, A., Kaviya, M., Balakrishnan, A., & Sridhar, T. (2023). Polymers and nanomaterials as gene delivery systems. In *Advances in Biomedical Polymers and Composites* (pp. 513-539). Elsevier.

Vishwakarma, S. K., Bardia, A., Tiwari, S. K., Paspala, S. A., & Khan, A. A. (2014). Current concept in neural regeneration research: NSCs isolation, characterization and transplantation in various neurodegenerative diseases and stroke: A review. *Journal of Advanced Research*, 5(3), 277-294.

Wade, J. (2023). A Historical Review of Hematopoietic Stem Cell Transplantation: Developmental Advancements Then and Now. Senior Honors Theses.

Wainwright, S. P., Williams, C., Michael, M., Farsides, B., & Cribb, A. (2006). From bench to bedside? Biomedical scientists' expectations of stem cell science as a future therapy for diabetes. *Social science & medicine*, 63(8), 2052-2064.

Walia, B., Satija, N., Tripathi, R. P., & Gangenahalli, G. U. (2012). Induced pluripotent stem cells: fundamentals and applications of the reprogramming process and its ramifications on regenerative medicine. *Stem cell reviews and reports*, 8, 100-115.

Wei, B., & Lu, J. (2021). Characterization of tendon-derived stem cells and rescue tendon injury. *Stem cell reviews and reports*, 1-18.

Weichselbaum, R., Goebbels, R., & Lokich, J. (1980). Complications of Cancer Therapy. In *Clinical Cancer Medicine: Treatment Tactics* (pp. 325-354). Springer.

Xu, J., Li, Y., Xu, Y., Ding, C., Li, T., & Zhou, C. (2014). A simple and effective method for the isolation of inner cell mass samples from human blastocysts for gene expression analysis. *In Vitro Cellular & Developmental Biology-Animal*, 50, 232-236.

Yedigaryan, L., Martínez-Sarrà, E., Giacomazzi, G., Giarratana, N., van der Veer, B. K., Rotini, A., Querceto, S., Grosemans, H., Cortés-Calabuig, Á., & Salucci, S. (2022). Extracellular vesicle-derived miRNAs improve stem cell-based therapeutic approaches in muscle wasting conditions. *Frontiers in immunology*, 13, 977617.

Yefroyev, D. A., & Jin, S. (2022). Induced pluripotent stem cells for treatment of Alzheimer's and Parkinson's diseases. *Biomedicines*, 10(2), 208.

Zakrzewski, W., Dobrzyński, M., Szymonowicz, M., & Rybak, Z. (2019). Stem cells: past, present, and future. *Stem cell research & therapy*, 10(1), 1-22.

Zhang, B., & Chen, T. (2024). Local and systemic mechanisms that control the hair follicle stem cell niche. *Nature reviews Molecular cell biology*, 25(2), 87-100.

Zhang, B., Stone, M. R. L., Sanjaya, K., Łapińska, U., Pagliara, S., & Blaskovich, M. A. (2022). Application of antibiotic-derived fluorescent probes to bacterial studies. In *Methods in Enzymology* (Vol. 665, pp. 1-28). Elsevier.

Zhang, X., Zhang, S., & Wang, T. (2022). How the mechanical microenvironment of stem cell growth affects their differentiation: A review. *Stem cell research & therapy*, 13(1), 415.

Zhang, Z. G., Buller, B., & Chopp, M. (2019). Exosomes—beyond stem cells for restorative therapy in stroke and neurological injury. *Nature Reviews Neurology*, 15(4), 193-203.

Zipori, D. (2005). The stem state: Plasticity is essential, whereas self-renewal and hierarchy are optional. *Stem Cells*, 23(6), 719-726.

About The Authors

Maria Nazir is an MPhil Scholar of Microbiology at the Cholistan University of Veterinary and Animal Sciences (CUVAS) in Bahawalpur, Pakistan. She is involved in a National Research Program for Universities (NRPU) Project, with research interests in Molecular Characterization. She has submitted abstracts to various conferences and authored and coauthored of several book chapters. Her research focuses on the isolation and Molecular characterization of *Corynebacterium pseudotuberculosis* and their applications in promoting animal health. Her interested research criteria are based on genetics, molecular biology, and bioinformatics.

Email: nazirmaria545@gmail.com

ORCID 0009-0004-6714-6212

Mr. Hameer Khan Khaskheli received his Master's in 2023 from the University of Padova, Italy. He is currently a Research Assistant in the Department of Biomedical Sciences at the University of Padova, Italy. His research expertise spans Molecular Biology, Cancer Biology, Medical Immunology, Stem Cell Research, and Precision Medicine. His primary focus is the intricate regulation of protein phosphorylation by kinases and phosphatases - critical processes often dysregulated in cancer and other diseases. He has an extensive record of publications in prestigious national and international journals and has authored two book chapters with a reputable publisher.

Email: hameerkhan.khaskheli@studenti.unipd.it
9282

ORCID 0009-0002-9622-

Miss Ammara Hameed received her MS in Biochemistry from The Islamia University of Bahawalpur by securing a 3.87/4.00 CGPA, in Punjab. She is serving as a biology lecturer at Army Public School and College Bahawalpur. Her research interest criteria are based on genetics, molecular biology, and bioinformatics. The research title is " Detection of NBS1 Mutations in Leukemia and Lymphoma Patients". She has reported two NBS1 novel mutations in Leukemia and Lymphoma Patients in Pakistan. She has two book chapters and one research article related to breast cancer under process.

Gmail ID: ammarahameed09@gmail.com

ORCID: 0009-0002-7337-2692

Mr. Tasawar Abbas received his M. Phil in Molecular Biology from the National Center for Excellence in Molecular Biology University of Punjab Lahore Pakistan. He has one review article. He has worked in Forensic Serology, He has excellent knowledge of Genetics, STRs, genetic polymorphism also Stem cells, and virology. His research interests include stem cell therapies, Immunology, Human genetics and criminology.

Email: tasawarabbas333@yahoo.com

ORCID: 0009-0002-7718-2634

Dr. Danish Riaz received his PhD (Zoology) in 2018 from Government College University Faisalabad, Pakistan. He is serving as Assistant Professor of Zoology at the Department of Zoology, Division Science and Technology, University of Education Lahore Pakistan. His research interests include Biological Science with a specialization of animal biotechnology and Fisheries. He has published over 34 research articles in well-reputed national and international journals. He also has written book chapters.

Email: danish.riaz@ue.edu.pk

ORCID-0000-0002-6137-8993

Shafeeq Ur Rehman earned his Bachelor in Applied microbiology from Cholistan University of Veterinary and Animal Sciences (CUVAS) Bahawalpur Pakistan. Currently he is pursuing M. phil in Microbiology from Cholistan University of Veterinary and Animal Sciences (CUVAS) Bahawalpur Pakistan. He is also working as research assistant in a project titled as "Robust Molecular Detection of Silent Circulation of FMD and PPR Viruses in Cholistan as a Model for Disease-Free Zone". His research interest is in FMD and PPR Viruses, Mucosal Immunity, Virology, Molecular cell

biology, Cell culture, Microbial genetics and CRISPR case tools for combating viral diseases. He has published 7 articles and many book chapters in national and international journals.

Email: shafeequrrehmanbhutta@gmail.com

ORCID: 0000-0003-3571-8226

Dr. Muhammad SAFDAR earned his PhD in Molecular Biology and Genetics from Gaziantep University, Turkey. He is Lecturer in the Breeding and Genetics department at Cholistan University of Veterinary and Animal Sciences (CUVAS), Bahawalpur, Pakistan. His research interests are molecular genetics and genomics, nutrigenomics, nano-genomics, bioinformatics, biotechnology, and their applications. He has published more than 70 research articles in national and international journals. He has also written many book chapters as well as an edited book. He is an associate editor for international journals.

E-mail: msafdar@cuvas.edu.pk

ORCID: 0000 0002 3720 2090:

To Cite This Chapter

NAZIR M et al., (2024). Stem cell research: unlocking the potential for regenerative therapies. In *Current Studies in Health and Life Sciences 2024* (pp.212-237). ISRES Publishing.

The Never-Ending Battle: Strategies for Infectious Disease Control in The Face of Emerging Pathogens

Faizan ALI

Atif AHMED

Samra MAQSOOD

Minahal FATIMA

Minal HUSSAIN

Shafeeq Ur REHMAN

Muhammad SAFDAR

1. Introduction

Both natural processes, such as the development of pathogens over time, and human behavior and behaviors may cause newly emerging infectious illnesses. Natural processes include the evolution of pathogens over time. Population expansion, migration from rural regions to urban areas, international air travel, poverty, conflicts, and detrimental ecological changes as a result of economic development and land usage are some of the factors that have led to these changes. The transmission of infectious agents from animals to people is the cause of a significant number of newly discovered illnesses (Fong & Fong, 2017). The probability that humans may come into close contact with animal species that have the potential to be hosts of an infectious agent is certain to rise as the human population grows and extends into new geographical locations. It is simple to understand that this combination creates a significant danger to human health when that element is paired with increases in human density and mobility. This combination poses a hazard to human health. As a potential contributor to the growth of infectious illnesses, climate change is becoming an increasingly pressing reason for worry. Diseases can spread into new geographical regions when the temperature of the Earth increases, and various ecosystems are changed (Baker et al., 2022; Jones et al., 2008). Certain microorganisms, such as bacteria, viruses, and others, can evolve over time and acquire resistance to the medications used to treat illnesses caused by infections. Consequently, medicines that were beneficial in the past for the purpose of illness control are no longer suitable for this purpose. Vaccine coverage that is decreasing may lead to the resurgence of disease; hence, even when there is a vaccine that is both safe and effective, an increasing percentage of individuals opt not to be vaccinated against the illness (Baker et al., 2022; Trovato et al., 2020).

The plague in the fourteenth century, the Spanish flu of 1918, and, recently, the COVID-19 pandemic are examples of threats that have not disappeared but are constantly resembling one difference: they remain deadly. They underscore the need for constant alertness and creativity in how people address the spread of infectious diseases (Mack et al., 2007). Given the availability of new pathogens, conventional approaches to combating diseases do not suffice. One of the reasons why attempts to fight this disease proved so difficult is that various factors influenced its evolution, including changes in the environment, growing international travel, and mutations of the microbes that caused the disease (Levitt et al., 2010). Technological changes such as deforestation and climate change affect ecosystems in a way that the disease-causing pathogens seek new niches to thrive on. Globalization also helps in the speedy interconnectivity of people and the carriage of goods, helping to spread diseases. However, pathogens are not stateless entities; they change, and evolution may, on certain occasions, compromise their susceptibility to treatments (Flood et al., 2024).

Solving these problems is possible only through a complex set of measures. Healthcare, prevention, and early diagnosis programs are pivotal to preventing severe epidemics. It has been